

MANITOBA ENVIROTHON
PLANT ECOLOGY
PROVINCIAL RESOURCES

ACKNOWLEDGEMENTS

We would like to thank:

Olwyn Friesen (PhD Ecology) for writing, compiling, and editing this document.

Subject Experts and Editors:

Barbara Fuller (*Project Editor*, Chair of Test Writing and Education Committee)

Lindsey Andronak (*Soils*, Research Technician, Agriculture and Agri-Food Canada)

Jennifer Corvino (*Wildlife Ecology*, Senior Park Interpreter, Spruce Woods Provincial Park)

Cary Hamel (*Plant Ecology*, Director of Conservation, Nature Conservancy Canada)

Lee Hrenchuk (*Aquatic Ecology*, Biologist, IISD Experimental Lakes Area)

Justin Reid (*Integrated Watershed Management*, Manager, La Salle Redboine Conservation District)

Jacqueline Monteith (*Climate Change in the North*, Science Consultant, Frontier School Division)



SPONSORS



“Earth laughs in flowers”

Ralph Waldo Emerson

| | |
|---|-----------|
| Fundamentals of Plant Biology..... | 7 |
| What is a plant?..... | 7 |
| Taxonomy and terminology | 7 |
| Vascular whole-plant structure..... | 9 |
| Internal structure of a vascular stem | 11 |
| External structures of woody plants..... | 12 |
| Internal structure of a leaf | 12 |
| Types of Plants | 14 |
| Non-vascular Plants..... | 15 |
| Vascular Plants | 16 |
| Gymnosperms | 16 |
| Angiosperms | 17 |
| Functional Plant Groups | 19 |
| Photosynthesis | 20 |
| Location of plant photosynthesis | 21 |
| Process of Photosynthesis | 22 |
| Variation in Photosynthesis | 23 |
| Diversity, form, and function | 23 |
| Leaves | 23 |
| Stems..... | 25 |
| Roots | 26 |
| Wood | 26 |
| Nutrition | 27 |
| Plant growth, development, and reproduction | 27 |
| Root Growth | 28 |
| Seed-free plants..... | 28 |
| Vegetative reproduction..... | 29 |
| Flowers | 30 |
| Seed Germination..... | 31 |
| Introductory Ecology | 33 |
| Adaptation and Acclimation..... | 33 |
| Ecosystem Constituents and Trophic Ecology | 34 |
| Producers and Consumers | 34 |
| Decomposers | 36 |
| Nutrient Cycling | 36 |

| | |
|--|-----------|
| Plant Associations | 38 |
| Mycorrhizae | 38 |
| Soils | 39 |
| Pollinators..... | 39 |
| Seed Dispersal | 41 |
| Habitats | 42 |
| Ecoregions | 42 |
| Plant Communities..... | 45 |
| Stratification and Structure..... | 45 |
| Vertical stratification | 45 |
| Slopes and Elevation | 46 |
| Habitat types | 47 |
| Grasslands..... | 47 |
| Shrubland..... | 50 |
| Deserts..... | 51 |
| Tundra..... | 52 |
| Forests..... | 53 |
| Competition | 57 |
| Succession of plant communities..... | 58 |
| Community stability and equilibrium | 59 |
| Disturbance-dependent ecosystems | 60 |
| Ecotones and Edge effects | 63 |
| Plants and their Environment | 64 |
| Winter Adaptations | 64 |
| Shedding Leaves | 64 |
| Plant Growth and Shape..... | 65 |
| Freezing Resistance | 65 |
| Wind Adaptations..... | 66 |
| Stems and Leaves..... | 66 |
| Seed Dispersal | 66 |
| Size and Clustering | 66 |
| Plant Adaptations to Extreme Environments | 66 |
| Ecological groups of plants: hydrophyte, xerophyte, mesophyte, halophyte | 67 |
| Hydrophytes | 67 |

| | |
|--|-----------|
| Xerophytes | 69 |
| Common environmental problems | 70 |
| Foliage burn | 70 |
| Pollution..... | 70 |
| Lack of water | 71 |
| Shade..... | 72 |
| Humidity and temperature..... | 72 |
| Environmental Conservation | 72 |
| Pesticides and Fertilizers | 72 |
| Invasive Species..... | 73 |
| Deforestation problems | 85 |
| Loss of biodiversity | 86 |
| Endangered Species | 86 |
| Climate Change | 88 |
| Skills and methods in plant science | 90 |
| Vegetation characteristics | 90 |
| Pathogens and Disease Management | 90 |
| Forestry | 93 |
| Forestry Management..... | 93 |
| Forestry Measuring Techniques..... | 94 |
| Agriculture | 96 |
| Sustainable agriculture | 96 |
| Horticulture..... | 97 |
| References..... | 98 |



FUNDAMENTALS OF PLANT BIOLOGY


Plants are fundamental to life on Earth. Plants support all aspects of human life — from our food, to our clothes, raw materials we use, oxygen we breathe, the medicines we take and much more. Essential services that plants provide are often taken for granted.

WHAT IS A PLANT?

Plants are any one of an immense number of species within the biological realm. Generally, plants are organisms that are considered to have limited motility and for the most part produce their own food. They include a number of recognizable species, including trees, shrubs, grasses, vines, ferns, algae, sea weed, and mosses. Plants are multicellular with cell walls containing cellulose. Plants are organisms that are capable of photosynthesis.

Taxonomy and terminology

Species have been traditionally divided into two kingdoms: plants and animals. In the age of DNA analysis and other techniques, fungi and bacteria have now been moved to form their own kingdoms. For example, research has shown that fungi have cell walls that use chitin instead of cellulose, as seen in plants. Lichen are unique as they are a symbiotic association between a photosynthetic organism, such as



algae or cyanobacterium and fungi, and so they are not considered plants.

The study of plants is often referred to as botany. As of 2017, scientists have identified approximately 391 000 species, within 452 groups, of vascular plants (a plant that has conducting tissue). Approximately 2000 new plants are identified annually (State of the World's Plants 2017). Of these identified plants, about 369,000 species (or 94 percent) are flowering plants. Further, more than 80% of the food derived from plants comes from 17 plant groups or families.

Scientists group and classify plants based on similarities. Plants are characterized by their morphology and DNA. Family is one such rank of classification that connects species that share some set of what are considered to be the more important or significant features. For example, vascular plants have lignified tissue and specialized structures termed xylem and phloem, which transport water, minerals, and nutrients upward from the roots and return sugars and other photosynthetic products. Vascular plants include ferns, club mosses, flowering plants, conifers and other gymnosperms.

Green plants – green algae and land plants – form a group consisting of all the descendants of a common ancestor. With a few exceptions, the green plants have the following common features:

1. Primary chloroplasts (organelle in plant cell) containing chlorophylls a and b,
2. Cell walls containing cellulose
3. Food stores in the form of starch contained within the plastids (organelle)
4. Undergo closed mitosis (cell replication) without centrioles
5. Mitochondria (organelle) with flat cristae
6. Chloroplasts (organelle) of green plants are surrounded by two membranes

There are two main groups of plants – non-vascular and vascular. Non-vascular plants include algae and bryophytes. Vascular plants are more numerous and include lycopods, ferns and horsetails, and the most common, gymnosperms and angiosperms. We have described and included a description of many of the plant groups below.

VASCULAR WHOLE-PLANT STRUCTURE

Vascular plants have three basic organs, roots, stems, and leaves.

Roots anchor the plant into the soil, absorb water and minerals, and store food. Roots have root hairs.

Taproot/Fibrous: There are two different kinds of root systems. Fibrous roots have many slender roots about the same size that spread out. Grass has fibrous roots. Taproots grow straight down, some as far down as 4.5m. Carrots and radishes have taproots.

Lateral/Secondary roots: these roots extend horizontally from the primary roots and anchor the plant into the soil. This branching contributes to water intake and facilitates the uptake of nutrients from the soil that the plant needs for growth.

The shoot system includes both stems and leaves:

Stems are the plant axis that contain the buds and shoots with leaves, and at their basal end, the roots of the plant. The stems also conduct water, minerals, and food to other parts of the plant. They also can store food and green stems can be locations where photosynthesis takes place.

Terminal bud: a bud that terminates the end of a stem or a twig

Bud: A young shoot from which leaves or flowers may develop

Lateral/Axillary Bud: A bud arising from a leaf.

Vegetative shoot: consists of stems and leaves


Node: critical areas from which leaves, branches, and roots grow out of the stem

Internode: areas between the nodes

Leaves are the flattened green outgrowth from the stem. The leaf is the primary site of photosynthesis, which produces food for plants. Leaves can be highly modified and adapted to a plant's surroundings. For example, the sharp spines of cacti, the needles of pines and other conifers, and the scales of an asparagus stalk.

Blade: broad, flat part of the leaf. A majority of photosynthesis occurs in the blade. The type of edge, pattern of veins, and number of blades per leaf can all be variable.

Petiole: this is the stem-like part of the leaf that joins the blade to the stem. Tiny tubes in the petiole connect the veins to the blade. Some of the tubes also carry water into the leaf, while others carry away food that the leaf has made. In some plants, the petioles can bend so that the blades can receive the most sunlight. The flexibility of the petiole also ensures the blade can twist in the wind and avoid damage.



The major parts of an angiosperm vascular plant. Only the parts of the plant discussed above are required to be memorized.

The reproductive systems are varied and are generally specific to a particular group of plants, such as flowers and seeds for flowering plants, sori for ferns, and capsules for mosses (see plant reproduction for more detail).

Internal structure of a vascular stem

Xylem: the xylem is one of two types of transport tissue found in vascular plants. The function of this transport tissue is to transport water and nutrients from the roots to shoots and leaves.


Phloem: the phloem is the other type of transport tissue found in vascular plants. The function of this tissue is to transport organic compounds made during photosynthesis, such as the sugar sucrose, to the parts of the plant where they are needed. In trees the phloem is the innermost layer of the bark.


Vascular Cambium: this is a layer of actively dividing cells, located between the xylem and phloem. This layer of cells is responsible for the secondary growth (increasing thickness) of stems and growth.

Stem: the stem is the plant axis that bears the buds and shoots with leaves and roots. The stem conducts water, minerals, and food to other parts of the plant.

Vascular bundles: the vascular bundle is part of the transport system in vascular plants. The vascular bundle is a strand of conducting vessels in the stem or leaves of the plant, containing the phloem and xylem.

Epidermis: the outermost layer of cells covering the stem, root, leaf, flower, fruit, and seed parts of a plant. At a particular part of the life cycle of *woody plants*, they stop growing in length and begin to grow in girth (width). The plant does this by growing secondary tissue around the circumference of the plant. The secondary tissue comes from the vascular cambium. A second cambium called the **cork cambium** is the source of **periderm**, which is a protective tissue that replaces the epidermis.






External structures of woody plants

Cork cambium: in woody plants comes arises near the surface of the plant. It produces cork cells on the outside. The cork cambium, cork cells, and phellogen form the periderm.

Periderm: protective tissue on the outside of the plant body. At maturity, the cork cells are non-living and the inner walls are lined with suberin, a fatty substance that gasses and water cannot penetrate.

Bark is another term used to refer to the combination of cork, cork cambium, periderm, and secondary phloem.



Internal structure of a leaf

Epidermis: the outermost layer of cells covering the leaf. With the cuticle, the epidermis provides a protective barrier against infection, water loss, and physical injury.

Stoma: microscopic openings on the surface of leaves and stems. Generally, there are more stoma on the underside of leaves. They function to allow the exchange of gases between the plant and the outside world.

Bundle sheath: a layer of cells in the leaves and stems of plants that form a sheath surrounding vascular bundles (containing phloem and xylem)

Cuticle: protective fil covering the surface of the leaf. It is made from lipids (fats) and polymers with wax.


Guard cells: these are cells that surround each stoma. They help regulate the transpiration rate (rate of taking in moisture) and gas exchange by opening and closing the stoma.

Mesophyll: this is the inner tissue of a leaf.

Spongy mesophyll: this layer of cells in the interior of a leaves consists of loosely arranged, irregularly shaped cells that have chloroplasts. This is the location of gas exchange for photosynthesis and respiration.

Palisade mesophyll: one or more layers of cells located directly under the upper epidermis. These cells are extremely important in photosynthesis.

TYPES OF PLANTS



Basic groupings of plants, starting from the most inclusive groupings (e.g., land plants), to the most exclusive groups (e.g., seed plants)

© Royal Botanic Gardens 2017

Non-vascular Plants

Green Algae

Chlorophyta: green algae, with members of this group all having motile swimming cells. Most species in this group can be found in aquatic environments, both freshwater and marine, a few species have adapted to a range of terrestrial, or land environments. For example, *Chlamydomonas nivalis*, which causes watermelon snow, lives on summer alpine snowfields.

Charophyta: freshwater green algae, often known as stoneworts or brittleworts. They are mostly found in slow-moving or standing water. These plants generally grow anchored to the surface by rhizoids.

Bryophytes

Bryophytes are small, non-vascular plants, such as mosses, liverworts, and hornworts. Over 12 000 species of moss are known to exist. Bryophytes play an important role in regulating ecosystems as they provide a buffer system for other plants, and these plants also benefit from the water and nutrients that bryophytes can collect.


Bryophyta: moss species; these are small flowerless plants that are mostly found growing in dense green clumps or mats, with wiry stems. Mosses often are found in damp or shady locations. Mosses reproduce by spores and they are considered to be non-vascular plants (lack of conducting tissues).

Hornworts: these are a group of non-vascular plants (lack of conducting tissues) that have a long horn like structure, called the sporophyte. Hornworts are found in damp or humid locations. They are characterized by each of their cells only have one chloroplast. Like mosses, hornworts reproduce using spores.



Watermelon Snow

© Royal Botanic Gardens 2017



Stoneworts

© Dave Carlson



Moss



Hornworts

Liverworts: these are a group of small non-vascular plants that most often grow in a form much like a flattened moss. They have single-celled rhizoids. They are most often found in humid locations, however, there are a few desert and Arctic species.



Umbrella liverwort

Vascular Plants

Seed-free vascular plants

They produce free spores, the principal dispersal units, via meiosis. Spore: a reproductive cell, capable of developing into an adult without fusion with another cell. Spores develop within a sporangium.

Lycopods: These species differ from all other vascular plants in having microphylls, leaves that have only a single vein (vascular trace) rather than the much more complex vein system found in ferns and seed plants. Plants from this group were used as flash early in photography and experimental photocopying machines as they have oil compounds in the cell walls that light on fire quickly and produce a flash of light. Today, many species of this group can be over-collected for Christmas wreaths. Lycopods include clubmosses, scale trees, and spike mosses.



Lycopod

Ferns and Horsetails: vascular plants lacking flowers and seeds, reproducing by spores. Horsetails are primarily colonizers of unforested areas, lake margins, and wetland. Leaves of horsetails are found in whorls and united to form a sheath around the stem. Ferns have leaves that are considered ‘megaphylls’, or leaf with several or many large veins branching apart or running parallel and connected by a network of smaller veins. Ferns fold their leaves into a bud, forming a fiddlehead, for protection.



Fern (fiddlehead)

Gymnosperms

Gymnosperms are a group of seed producing plants that includes conifers, cycads, *Ginkgo*, and gnetophytes. Gymnosperm seeds develop on the surface of scales or leaves, which are often modified to form cones.

Pinophyta: more commonly known as conifers, this group of plants have a vascular system and cones (containing seeds), but no flowers. Most conifers are trees, although a few are shrubs. Examples include cedars, Douglas firs, cypresses, firs, junipers, kauri, larches, pines, hemlocks, redwoods, spruces, and yews. It has been estimated that there are over 820 species in this group.



White Pine



Cycad







Ginkgo

Angiosperms

Angiosperms are also known as the flowering plants. There are over 300 000 known species of flowering plants and remain the dominant group of land plants. Flowers within this group have remarkable variation in form and elaboration. Most of the fruits and vegetables humans consume come from this group. Over 80% of the food on earth comes from 17 plant families found within this group. The cotton and linen in our clothing is also from angiosperms. Four features define angiosperms: (1) possession of flowers (with stamens and ovaries (which become fruit!)), (2) double fertilization (two nuclei in plant sperm), (3) less reproductive stages, and (4) specialized tissues for transporting water, known as the vessel elements.

Monocots: large group of plants that includes familiar plants such as grasses, lilies, irises, orchids, cattails, and palms. This large group of plants is characterized by trimerous flowers and usually parallel-veined leaves.

Eudicots: extremely large group of plants (over 175 000 species), including most familiar trees and shrubs (e.g., oaks and apple trees), as well as many herbs (e.g., sunflowers, petunia, and buttercups). They are characterized by 4- or 5-merous flowers, and usually branching-veined leaves.

| | Seed | Root | Stem | Leaf | Flower |
|----------|---|---|--|---|---|
| Monocots |  One cotyledon in seed |  |  |  |  |
| Eudicots |  Two cotyledons in seed |  |  |  |  |

© The McGraw-Hill Companies, Inc.

Magnoliids: smaller group of plants (about 9,000 species), including magnolias, nutmeg, bay laurel, cinnamon, avocado, and black pepper. They are characterized by trimerous flowers, pollen with one pore, and usually branching-veined leaves.



Magnoliids

Functional Plant Groups

We often use general names to define groups of plants with some similarities in function or appearance.

Graminoid: graminoid refers to an herbaceous plant with a grass-like morphology, i.e. elongated culms with long, blade-like leaves. They are contrasted to forbs, herbaceous plants without grass-like features.



Graminoid and grasses

Grasses: Most grasses have round stems that are hollow between the joints, bladelike leaves, and extensively branching fibrous root systems. Grasses are any of many low, green, nonwoody plants belonging to the grass family (Poaceae), the sedge family (Cyperaceae), and the rush family (Juncaceae). There are many grass-like members of other flowering plant families.

Forbs: A forb is an herbaceous flowering plant that is not a graminoid (grasses, sedges and rushes).

Herbs/Herbaceous: are plants that have no persistent woody stem above ground.

Trees: a woody plant with an elongated stem, or trunk, supporting branches, and leaves in some species. Trees can include both angiosperm and gymnosperm species.

Epiphytes: organisms that grow on the surface of plants and obtain moisture and nutrients from the air, rain, water, or debris around it. Epiphytes use the plants they grow on as support but do not necessarily negatively affect the plant. Mosses, algae, liverworts, orchids, and bromeliads are examples of plant epiphytes.

Annual plant: these are plants that completely die off at the end of a growing season or when they have flowered or fruited. The need to grow again from the seeds they have produced.



Sunflower: Forbs



Trees

© Ed Yourdon



Fox Brush Orchid: Epiphytes

© Dinesh Valke


Perennial: these are plants that have stems that die at the end of the season, but parts of the plant survive from season to season. The surviving part of the plant can include roots, a caudex (thickened portion of the stem at ground level), bulbs, and tubers.



Japanese peony: Perennial

PHOTOSYNTHESIS

Photosynthesis is the process by which green plants and certain organisms use solar energy to convert water and carbon dioxide into a simple sugar, **glucose**. As such, photosynthesis provides the basic energy source for nearly all organisms. Further, the process of photosynthesis creates oxygen as a by-product, an essential element that most organisms depend on.



Photosynthesis occurs in green plants, algae, and some species of bacteria. These plants produce millions of new glucose molecules every second. Plants will use this glucose, which is a carbohydrate, to build their leaves, seeds, flowers, and fruit. They are also able to take this glucose and convert it to cellulose. **Cellulose** is used as a structural material in cell walls. When plants produce more glucose than they can use they can store it in the form of starch and other carbohydrates in their roots, stems, and leaves. If conditions change, plants can use this excess glucose for extra energy or building materials.

Photosynthesis is a key process in the functioning of all ecosystems. Humans and other animals depend on glucose as an energy source but are unable to make it. This creates a reliance on plants as our primary source of energy. Further, the oxygen produced as a by-


product of photosynthesis is critical for all humans and other animals. Finally, humans also depend on the ancient products of photosynthesis, known as fossil fuels. The complex mix of hydrocarbons that make up fossil fuels are the remains of organisms that relied on photosynthesis millions of years ago. Thus, all life on earth depends on photosynthesis, either directly or indirectly, making it the most important biochemical process.

Photosynthesis relies on the photosynthetic pigment **chlorophyll** to complete this process. During the light reactions of photosynthesis, as detailed below, solar energy or light energy excites electrons in chlorophyll and boosts them to a higher energy level. This energy is used to drive the production of glucose. The chlorophyll also gives leaves and plants their distinct green colour. In the autumn, as days get shorter, leaves of certain plants (e.g., deciduous trees) begin to die and the chlorophyll degrades and disappears from the leaves. The lack of chlorophyll allows the other photosynthetic pigments found in the leaf to be seen, which gives us the characteristic autumn colours.

Location of plant photosynthesis

Photosynthesis occurs in the green stems and leaves of plants. Each plant has millions of cells with specialized cell structures known as chloroplasts. Each cell in a plant leaf contains 40-50 chloroplasts. The chloroplast is an oval-shaped structure that is divided into compartments by membranes. Each disc shaped compartment, known as a **thylakoid**, is arranged vertically within the chloroplast like a stack of plates. Each stack of thylakoid's is known as a **grana**, and they are suspended in a fluid known as **stroma**.

Chlorophyll, a light-trapping pigment, is found in the **membranes** (a pliable sheet like structure acting as a boundary, lining, or partition) of thylakoids. All the other pigments, **enzymes** (organic substances that speed up chemical reactions), and other molecules needed for photosynthesis are also located in these membranes.



Structure of a plant cell. Notice the cell wall and chloroplast. Only the parts of the plant cell discussed in the text are required to be memorized.

Process of Photosynthesis

Photosynthesis is a complex process. Plant biologists divide it into two stages:

Stage 1 – Light Dependent Reaction

Chloroplast trap light energy and convert it into chemical energy (NADPH and ATP).

Photosynthesis begins when light strikes pigments and excites their electrons. The energy becomes absorbed (by P700 and P680) and pass through a chain of carrier molecules (electron transport chain). Some of the energy then combines with a hydrogen ion and NADP⁺ to form NADPH. Some of the energy gets used to produce ATP. Finally, some of the energy is used to split water into oxygen, hydrogen ions, and electrons. Oxygen from the water get released through the leaf.

Stage 2 – Light Independent Reaction

This stage is also known as the Calvin cycle. Chemical energy (NADPH and ATP) provide the energy for the reactions used to synthesize glucose. These reactions occur in the stroma. The Calvin cycle is dependent on the presence of carbon dioxide molecules, that enter the

plant through pores in the leaves. The light-independent reaction begins when carbon dioxide molecules link to sugar molecules known as RuBP. With an enzyme, through several intermediate steps, glucose is produced.

Variation in Photosynthesis

Environmental conditions can impact how plants carry out photosynthesis. Plants such as corn and crabgrass have evolved to live in hot, dry environments and must overcome certain obstacles to photosynthesis. These plants partially close the pores of their leaves on hot days to reduce the amount of water escaping. When the plants partially close their pores, this reduces the amount of carbon dioxide that can enter the leaf and pauses photosynthesis. To solve this problem, some hot-weather plants stockpile carbon dioxide in the stroma and slowly release the chemical to be able to continue to photosynthesize. This adaptation can allow plants to continue to photosynthesize on very hot days even when their pores are completely closed.

DIVERSITY, FORM, AND FUNCTION


Leaves

Edges: the type of edges on leaves can be varied. Almost all narrow, grass like leaves and needles leaves have a blade with a **smooth edge**, as do many broadleaf plants, particularly those that are native to warm climates. The rubber plant is a good example of such a plant.

Temperate broadleaf plants often have small, jagged points called **teeth** along their blade edge (or **serrations**). Birch and elm trees are great examples of this type of leaf.

Hydathodes, or tiny valve like structures, can be found on some leaves to release extra water. In the leaves of other plants, like cottonwood and pin cherry trees, there are special glands that produce liquids that protect the young leaf from being eaten by plant-eating insects.

Other trees, like the mulberry and oak trees, have **lobed** leaves. The leaves appear as if large bites have been taken out. This shape allows heat to leave the leaf.



Veins: Differences in veins can be found throughout vascular plants. Veins carry food and water, as well as support the blade. In many broad leaves, the veins can form a pattern like a net, with several large veins connected to small veins. The smallest veins will supply water to every part of the blade. They will also collect all of the glucose made by the leaf.



We often group the types of **net veins** into two groups, **pinnate** (feather like) and **palmate** (hand like). Leaves that are pinnate have one large central vein, **midrib**, which extends from the tip of the blade to the base.

Other veins branch off of each side of the midrib. Beech, birch, and elm trees all have this vein pattern. Palmate leaves have several veins that are equal in size that all extend from the base of the blade. Maple, sweet gum, and sycamore trees all have palmate leaves.

Needle leaves and narrow leaves do not have net veins. Narrow leaves have parallel vein patterns. Several large veins run alongside each other from the tip to the base of the blade. The large veins are connected by small cross veins. Needle leaves are much smaller, so they only have one or two veins running through the centre of their blades.

Blades: The number of blades per leaf and their arrangement vary between plants and can be used to classify and identify plants. A **simple leaf** is a leaf with only one blade. Grasses, apple and oak trees, and many other plants have simple leaves. A **compound leaf** has more than one blade. Each of the blades of this compound leaf are called **leaflets**.

Leaflets in a compound leaf can be arranged in a **pinnate** or **palmate** pattern. Pinnately compound leaves have leaflets growing in two rows, one on each side of the **rachis**, or central stalk. Garden peas, ash and walnut trees all have pinnate leaves. Leaves




that are palmate have leaflets that all grow from the tip of the leafstalk. Plants like clover, horse chestnut, and others have palmate compound leaves.

Double compound leaves can be found on a few plants, like carrots and Kentucky coffee trees. Each leaflet is divided into a number of still smaller leaflets on a double compound leaf. These leaves look more like a group of leaves and twigs than a single leaf.


Leaf Arrangements: Leaves can be arranged differently around the stem. In the **opposite** arrangement, there will be two leaves at a point of attachment (node) on opposite sides of the stem.

In the **alternate** arrangement, there will be one leave per point of attachment with the second leaf being above the first but attached to the opposite side of the stem.

Finally, in the **whorled** arrangement there are three or more leaves attached at one node.




Stems



Modified stems can be found on some plants and serve a variety of functions. For example, strawberries have modified stems called **stolons** that grow on the surface of the ground. They allow strawberries to spread and occupy a large area of nutrient-rich soil. **Tubers**, the modified stems of potatoes and other plants, are adapted for food storage. Some plants have **bulbs** that are modified stems for storage. **Rhizomes** are modified stems that grow laterally underground, and they are often mistaken for roots. The rhizome is used to store starches, proteins, and other nutrients. Finally, **corms** are short, vertical, swollen underground plant stems that also serve as storage organs for plants to survive harsh conditions. Corms are easily confused with bulbs. Corms are stems that have solid tissues, whereas bulbs are made from modified leaves.

Roots

Some plants have adapted to their environment by modifying their roots. **Adventitious roots** grow from the stem above the ground. They include the **prop roots** of corn and other plants. **Prop roots** grow down into the soil and help the plant hold itself brace against the wind. **Aerial roots** of orchids and other plants cling to branches and absorb water and minerals from the surface of the tree and from the air. The roots of mistletoe are known as **sinkers** and penetrate the limbs of a tree. Sinker roots absorb food, water, and minerals from the tree.



Wood

Wood is a fibrous and porous structural tissue that can be found in the roots and stems of trees and other woody plants. Wood is an organic material and is composed of cellulose fibers that are embedded in a matrix of lignin. Within trees, wood functions as a support, allowing the plant to grow large and stand by themselves. Wood also conveys water and nutrients between the roots, leaves, and other growing tissues.

Hard and soft wood: Wood is often classified as soft or hard. Generally, wood from deciduous trees (e.g., oak) is considered hard, while wood from coniferous trees (e.g., pine) is considered soft. Hardwoods tend to be slower growing and therefore are usually denser. However, hardwood is not necessarily a harder material and vice versa.

Nutrition

Plants have a variety of essential nutrients necessary for growth and metabolism. Carbon, oxygen, and hydrogen are obtained from the air, whereas other nutrients, including nitrogen, are normally obtained from the soil. There are seventeen nutrients that are the most crucial for plants.

Macronutrients: nitrogen (N), phosphorus (P), potassium (K), calcium (Ca), sulfur (S), magnesium (Mg), carbon (C), oxygen (O), and hydrogen (H).

Micronutrients: iron (Fe), boron (B), chlorine (Cl), manganese (Mn), zinc (Zn), copper (Cu), molybdenum (Mo), and nickel (Ni).

Many of these nutrients stay in the soil as salts, so plants uptake these elements as ions. Hydrogen, oxygen, nitrogen, and carbon contribute to over 95% of the biomass of plants. Micronutrients only make up 0.02% of the plant. As plants so strongly depend on the availability of nutrients in the soil, the composition of the soil will dictate the type of plants that can thrive within a region. In agriculture and horticulture (practice of garden cultivation and management), the addition of extra nutrients through fertilizers, can assist in the growth and yield of plants.


PLANT GROWTH, DEVELOPMENT, AND REPRODUCTION

Plant growth is directed by environmental and ecological factors. Environmental factors include temperature, precipitation, wind speed and direction, sunlight, soil nutrients, soil moisture, soil granularity and compaction, as well as the topography of the growth area. Other ecological factors, including water competition, nutrient competition, and light availability also impact plant growth. Herbivory and trampling from animals within the ecosystem also play a role in how quickly plants grow and develop.

Root Growth

Roots grow from their **apex**, or tip. The area of growth is known as the **apical meristem**. The **meristem** is a part of a plant where cells are dividing rapidly, continually forming new cells. The root cap covers the apical meristem and protects the root tip from damage as it moves through the soil.


Apical meristem cells are identical and small. In the **zone of elongation** the cells quickly grow longer. Further back in the **zone of maturation**, cells differentiate, which means they take on a different structure and appearance based on their future function in the mature root.



Seed-free plants

Seed-free plants reproduce through the production of free **spores**, the principal dispersal units, via meiosis. A spore is the reproductive cell that is capable of developing into an adult without fusing with another cell. Spores develop in the **sporangium**. These spores

germinate and develop into gametophytes that can live separate from their parent plant. Like all plants, seed-free plants produce male and female gametophytes. The sperm and egg form a zygote through fertilization. The sporophyte then develops from the zygote into an adult plant.




Copyright © 2008 Pearson Education, Inc., publishing as Pearson Benjamin Cummings.

© Pearson Education, Inc.


Vegetative reproduction

Vegetative reproduction is a form of asexual reproduction where a new plant grows from a fragment of a parent plant or grows from a specialized reproductive structure. Although numerous plants reproduce by vegetative reproduction, it is rare that this is the only method by which they reproduce.



Flowers

Floral structure and the parts of a flower are very important in how we group species of plants together. Flowers are highly specialized shoots that are made from stems and leaves



Parts of a Flower

1. Pedicel (flower stem) – this is the stalk supporting the flower
2. Receptacle – modified floral stem that the modified leaves (petals) or floral appendages arise from
3. Sepal – the outermost ring of leaves (usually green) that protect the inner floral parts of the flower while the flower is a bud
4. Petal – the second ring of leaves (usually coloured to attract pollinators) that surround the internal portions of the flower
5. Stamen – the male structure of the flower, made up of the anther and the filament.
 - a. Filament – slender stalk of the stamen supporting the anther
 - b. Anther – portion of the stamen that contains the pollen grains (sperm)
 - c. Pistil – the female structure of the flower, sometimes also referred to as the carpel, comprised of the ovary, ovule, stigma, and style
 - d. Ovary – base portion of the pistil that contains ovules (with eggs). When the flower matures this becomes the fruit with seeds.

- e. Ovule – fertile portions of the pistil that contains female gametophytes (eggs in embryo sac) that develop into seeds after fertilization.
- f. Stigma – receptive portion of style that receives and recognizes pollen (male gametophyte)
- g. Style – slender stalk of pistil above the ovary that pollen tubes must pass to reach the eggs

Flowers usually have some level of symmetry. Flowers that are radially symmetrical are known as **actinomorphic** (ray-form). Flowers that are bilaterally symmetrical are known as **zygomorphic** (one plane of symmetry).



Actinomorphic flower




Zygomorphic flower

Seed Germination

Seeds are living entities. They have embryonic tissue inside that is needed in order to germinate. Most seeds also have a store of food reserves with the embryo that are all wrapped in a seed coat. Seeds will germinate, or ‘wake up’, when the soil moisture and temperature conditions are correct for them to grow. Each plant species’ seeds have individual needs.

Seed dormancy: some seeds may not germinate as they have developed a dormancy or sleep period. These seeds will not germinate even if conditions, soil moisture and temperature are ideal. Staggering germinating benefits seeds to




Development of a eudicot seedling
© W.H. Freeman and Company

make sure some seedlings develop later and may avoid bad weather or herbivores that may have eaten them if they had germinated sooner.

Steps of Seed Germination

1. **Imbibition** – seeds rapidly take up water causing the seed coat to swell and soften.
2. **Interim or lag phase** – the seed activates internal physiology, cells begin respiration, and the seed starts to make proteins and metabolizes the food stores found underneath its seed coat.
3. **Radicle and root emergence** – As cells develop, elongate, and divide, the **epicotyl** and **radicle** (primary root) emerge from the seed.



Early Seedling Development

The radicle (primary root) emerges from the seed and anchors the plant to the ground. This allows it to start absorbing water. As the root absorbs water the shoot emerges from the seed.

In **eudicots** (two-seed leaves – see plant classification) have three main parts: **cotyledons** (seed leaves), **hypocotyl** (shoot under leaves), and **epicotyl** (shoot above the leaves).

Generally, the shoot either emerges from the soil in a hook shape with the cotyledons and shoot tip being pulled into the air, or only the section above the cotyledons expands so the cotyledons (seed leaves) decompose underground.

In **monocots** (one-seed leaves – see plant classification), the primary root is protected by a sheath (coleorhiza), that pushes out of the seed first. The seedling leaves also emerge covered in a protective sheath.

After the shoot emerges, the seedling grows slowly while using up the energy stored in the seed. The seedling soon develops a branched root system or taproot. True leaves, like the leaves of the mature plant, appear allowing the plant to photosynthesize.


INTRODUCTORY ECOLOGY

Plants are the most important source of primary production in all ecosystems on Earth. They produce the bulk of biomass from carbon dioxide, basic nutrients, and light. The foundation of this primary productivity is photosynthesis. This process has fundamentally altered the atmosphere of Earth, resulting in air that is 21% oxygen, which is essential to other living organisms. Plants also provide a food source for animals, as well as shelter and nesting locations for many animals.

Ecology is the study of organisms and their environment – and the interrelationships between the two. The **ecoregion** is a community of animals and plants that have common characteristics for the environment they live in. **Ecosystems** are the complex of living organisms, their physical environment, and their interrelationships in a particular area. A **community** is the populations of all the species, e.g., moose, wolves, rabbits, owls, beavers, trees, grasses, shrubs, etc., that live in the same area at the same time. **Populations** are the group of organisms from the same species that can interbreed and live in the same place at the same time, like the elk or grass in the figure below. Finally, an **individual** is an organism, like the elk or a tree in the figure below.

Adaptation and Acclimation

Adaptation is any behavioural, morphological, or physiological trait that is a result of natural selection. This inherited characteristic should enhance an organism's ability to survive and reproduce in their environment. Some individuals, who often possess these adaptations, will leave more offspring than



others. These individuals are considered to be more ‘fit’ than others because they contribute the most to the entire population’s gene pool. Differences in the reproductive success of individual organisms come about through the process of natural selection. Under a specific set of environmental conditions, the individuals that survive the best or have adaptations to best survive those conditions are selected for. Any individuals that either do not have adaptations to survive and reproduce in these conditions or survive worse than others will be selected against.

Acclimation is the short-term response of an individual to different or changing natural environments. For many species, this acclimation occurs each season. For example, many plants, such as maple trees, irises, and tomatoes could survive freezing temperatures if the temperature gradually decreases over days to weeks. However, if the same drop would occur over a short period (e.g., one day) the difference in temperature could kill the plants. The short period did not provide enough time for the plant to acclimate to the colder temperatures. Acclimation is also reversible. If a plant gets used to colder temperatures in the fall, it can also become acclimated to the warmer conditions in the spring.

ECOSYSTEM CONSTITUENTS AND TROPHIC ECOLOGY

Generally, we consider there to be four constituents to an ecosystem. There are **abiotic components, producers, consumers, and decomposers and nutrient cycling**.

The abiotic components of an ecosystem are the essential nonliving elements. They can include the air, the water, temperature, and rocks and minerals that make up the soil. Components can also include how much rain or snow falls, if there is fresh or salt water, how much sun an area gets, and its temperature range. All the other biotic elements of the ecosystem interact and depend on these abiotic components.

Producers and Consumers

Trophic ecology is the study of how energy moves through an ecosystem. All organisms must obtain energy for their growth, survival, and reproduction. The methods of obtaining these resources and the impacts of resulting interactions are all studied within trophic ecology.

Autotrophs – organisms that use inorganic sources of carbon and energy from solar radiation. Examples include plants, algae, and certain bacteria. They are also known as **PRIMARY producers**.

Heterotrophs – organisms that use organic sources of carbon by consuming other organisms or their by-products. Examples include animals, bacteria, and fungi. They are often referred to as **SECONDARY producers**.

Consumers – these are heterotrophs that consume other organisms

Decomposers – these are heterotrophs that consume dead organic matter or waste products


Herbivores – organisms that primarily consume plant materials. They include grazers (feed on leafy material like grasses), browsers (feed on woody material), granivores (feed on seeds), and frugivores (feed on fruit).

Carnivores – organisms that are ‘flesh-eaters’. They consume herbivores or other carnivores. Individuals that feed directly on herbivores are considered first-level carnivores (second level consumers). Individuals that consume both herbivores and first-level carnivores can be considered second-level carnivores (third level consumers).

The trophic level is a stem in the transfer of energy, or food, within a food web or chain.

There may be several trophic levels within a system, including primary producers, primary consumers, and secondary consumers. Further carnivores may form fourth and fifth levels. Primary producers are the most abundant food source and biomass (mass of organic material) available. Primary consumers, who consume primary producers, are the second most abundant group of organisms.

Tertiary and quaternary consumers represent the smallest groups of organisms. The amount of energy in each a trophic level is reduced with every step up.



Decomposers

Decomposers are living organisms that break down dead organisms and waste materials. Earthworms, dung beetles, nematodes, and many species of fungi and bacteria are all decomposers. These species form a crucial function within an ecosystem, as they recycle nutrients, returning them into the soil where plants can take them up again.




Fungi (decomposer) growing on a tree

Plants play an important role in water and nutrient cycling within an ecosystem. Some plants have coevolved with nitrogen-fixing bacteria, making plants an important part of the nitrogen cycle.

Nutrients move through ecosystems in biogeochemical cycles. These cycles include chemical exchanges of elements, such as carbon, nitrogen, and phosphorus, through the atmosphere, rocks, water, and living species, including plants and animals. As seen in the figure below, plants can play a critical role in the movement of nutrients within an ecosystem. The primary productivity of an ecosystem depends on plants taking up essential mineral nutrients (e.g., nitrogen, carbon, phosphorus), and incorporating these nutrients into their tissues, such as leaves and stems.


The nutrient cycling in all ecosystems involve three major components of inputs, internal cycling, and outputs.

The **carbon cycle** is a very important biochemical cycle as all organic compounds contain carbon and carbon is an essential for photosynthesis. The carbon cycle includes inputs of carbon dioxide from the atmosphere and dissolved in water. The carbon is internally cycled through animals and plants through photosynthesis (plants) and consumption (animals). Some carbon is also released (output) by plants and animals through respiration (e.g., humans breathing). Decomposing organisms, as well as factor emissions are considered outputs, releasing carbon dioxide back into the atmosphere and water.




A schematic drawing of nutrient cycling through a tree.

Modified from Smith and Smith 2001



Nitrogen cycles include an *input* of atmospheric nitrogen that gets taken up by nitrogen-fixing bacteria. These bacteria can often be found living on root nodules of some plants or within the soil. The bacteria convert the nitrogen to a form that can be assimilated by plants and *internally cycled* between plants and animals. Animals, and decomposers, including bacteria and fungi, release ammonium that can be converted to nitrates (by bacteria) and this *output* is eventually released back into the atmosphere.




PLANT ASSOCIATIONS

Plant roots play an essential role in soil development and prevention of soil erosion.


Mycorrhizae

At least 80% of vascular plants form mutually beneficial associations, known as mycorrhizae, between their roots and fungi. These play a key role in nutrient cycling and plant nutrition. Mycorrhizal fungi benefit their host plants by increasing its ability to capture water and essential elements, especially phosphorus. This type of fungi can also increase the absorption of zinc, manganese, and copper, three other essential nutrients. Mycorrhizae can also help protect the plant from pathogenic fungi and nematodes (small animals). In return for this assistance to the plant, the mycorrhizae receive carbohydrates and vitamins from the plant, essential to their growth.



Endomycorrhizae – mycorrhizae that **penetrates** the root cells. The most common of the mycorrhizae and is found in 80% of all vascular plants. The hyphae of the fungi move into the root cells.

Ectomycorrhizae – mycorrhizae that **surrounds** the root cells. These types of mycorrhizae are mostly found in certain groups of trees and shrubs in temperate regions, such as oaks, willow, poplar, birch, and pine. These associations help the trees be more resistant to the harsh, cold, and dry conditions that could limit their growth.



Soils

Soils consist of layers called horizons. As you examine a vertical section of soil, one can see variations in colour that is related to the amount of living and dead organic material, to porosity (porosity), the structure, and the extent of weathering. These differences result in a succession of areas, known as horizons. Generally, these horizons are labeled from A-C, with horizon A representing the topsoil, B representing the subsoil, and C representing the soil base which sits atop the bedrock. The pore space within the soil contains air and water, which are crucial for plant growth and development.


The top region, the **A horizon**, is often also known as the surface or top soil. In this uppermost region, the greatest physical, chemical, and biological activity is occurring. It is the layer where **humus** (dark coloured mixture of organic decaying products) accumulates. This layer is full of roots, insects, other arthropods, nematodes, earthworms, protists, nematodes, and other decomposing organisms.

The middle region, the **B horizon**, is also known as the subsoil. It is known as the region of deposition, containing substances such as iron oxide, clay particles, and small amounts of organic materials from the horizon above. This horizon can contain roots but contains much less organic material than the horizon above.

The final horizon, the **C horizon**, contains broken down and weathered rocks and minerals which help form the true soil in the horizons above.

Pollinators

Pollination is one of the essential processes of plant life. Pollen, which is from the stamen (male portion of the flower), is transported to the pistil (female portion of the flower). This allows a plant to produce seeds that become the next generation. The movement of this



pollen often requires outside influence, such as wind or animals. **Pollinators** are animals that help move the pollen.

Bees, butterflies, and moths are often what we think of when we think of pollinators. However, flies, beetles, wasps, ants, hummingbirds, and in some areas bats, are also all very important pollinators.


The pollen produced by flowering plants can also serve as a source of protein for many species of insects. Many of these plants also produce nectar, which is a sugar-based high energy food. Both nectar and pollen are attractive to pollinators. Many move from flower to flower, feeding and collecting pollen and nectar. This movement unintentionally transfers pollen from one flower to another and is known as **cross-pollination**. Genetic diversity and resilience are ensured by this cross-pollination.

Many flowering plants and pollinators have co-evolved (evolved together) to ensure the correct pollinator feeds from the correct plant. As seen in the figure, a flower wants the pollinator to spread its pollen to other flowers of the same species. As the flower evolves to be deeper, the fly responds and evolves a longer proboscis (mouth part) to match this change and access the pollen and nectar.

Plant-pollinator coevolution

Host specificity drives coevolution


A flower “wants” its pollen spread to other flowers of the same species



A coevolutionary escalation

Extravagant traits can coevolve in response

Flower evolves
Fly responds



Seed Dispersal

Plants that have heavy seeds that cannot be distributed by the wind will often use animals to carry their seeds away from the parent plant. This allows the seeds to be distributed in areas that may be better for their development and growth, as well as spread a plant throughout a habitat. Some animals that do this distribution also consume the seeds for their own nutrition. Plants that rely on this type of seed distribution need to produce an enormous number of seeds within their lifetimes as few of the seeds will survive, be deposited at a suitable site, and germinate.



The relationship between the Clark's nutcracker and the whitebark pine is one such animal plant relationship. Whitebark pine produce large wingless seeds that can be dispersed by animals. The seeds are eaten and hoarded by several animals, including chipmunks, Steller's jays, and Clark nutcrackers. However, only the Clark nutcrackers can appropriately disperse the seeds systematically and successfully away from the parent tree. The nutcracker carries the seed in cheek pouches and caches (stores) the seeds deep enough in the soil of the forests and fields to avoid them being eaten by other animals. An individual nutcracker can store around 98 800 seeds per year! The nutcrackers only eat a number of these seeds, which allow those forgotten or ignored to germinate in the soil and produce new trees. Whitebark pines are so dependent on Clark's nutcrackers to distribute their seeds that these two species, plant and animal, are almost always found within the same region.



Some plants enclose their seeds in a nutritious fruit that is attractive to fruit-eating animals. Fruit-eating animals, or frugivores, do not eat the seed but they eat the endocarp, or nutritious fruit, that surrounds the seed. Plants make sure these fruits are not eaten too early, before the seed is mature, by cryptic colouration, such as a green fruit among green leaves, and by making the fruit have a bad texture, and sometimes a hard outer coat. When the seed is mature, plants will attract animals by presenting attractive odours, changing the fruit texture, increasing the sugar and oils in the fruit, and through colour to catch attention.

HABITATS

A **habitat** is the place where an organism or community of organisms live. It includes all the living (biotic) and non-living (abiotic) conditions that an organism needs to survive. A **microhabitat** is the conditions and organisms in the immediate area of a plant or animal.


Abiotic factors, such as temperature, water, sunlight, wind, rocks and soil, and climate all impact a plant's ability to obtain the resources they need to live and ability to survive in the environment. The temperature of an area affects all biological processes. The availability of water within regions affects species distribution as all species need water to survive and many species live within this water. Sunlight provides the energy that plants use to grow. As the primary food source, the abundance and distribution of plants in an environment will impact the abundance, density, and diversity of plants in a region. Additionally, the physical structure of rocks and soil limit the distribution of plants and thus the animals that rely on them.

Climate is one of the biggest abiotic driving factors that influence the distribution of wildlife on earth. Climate influences the temperature of a region, availability of water, sunlight, and wind, as well as the structure of rocks and soil. It also limits the biological process of all living organisms and thus plays a large role in dictating the diversity and abundance of plants.

Plants have adapted to the habitat in which they live. Specialized adaptation of each species to their habitat ensures their survival and continued ability to reproduce. This adaptation also allows species to survive predictable changes in their environment, such as the onset of winter or summer, or the wet or dry seasons.

ECOREGIONS

Ecoregions are major ecosystems that result from a combination of predictable patterns of climate, which are influenced by latitude, global position, and altitude. Every ecoregion is a geographical area across which the interactions between climate, soil, and topography are uniform enough to allow for the development of similar types or



© 2014 R.G. Bailey/Springer Science

forms of vegetation. Ecoregions occur in predictable locations in different parts of the world.

Within ecoregions, scientist use a hierarchy to describe ecosystems. A **domain** represents a subcontinental area of related climates; there are four domains, polar, humid temperate, humid tropical, and dry. A **division** represents a single regional climate. Currently, 14 divisions are recognized (e.g., tundra, subarctic, subtropical, prairie, etc.).

| Domain | Division | Temperature | Rainfall | Vegetation |
|------------------------|------------------|---|--|--|
| <i>Polar</i> | Tundra | Mean temperatures of warmest month $<10^{\circ}\text{C}$ | Water low during the cold season | Moss, grasses, and small shrubs |
| | Subarctic | Mean temperature of summer is 10°C , winter - 3°C | Rain even throughout year | Forest, parklands |
| <i>Humid temperate</i> | Warm continental | Coldest month below 0°C | Adequate throughout year | Seasonal forests, mixed coniferous-deciduous forests |
| | Hot continental | Coldest month below 0°C , warmest month $>22^{\circ}\text{C}$ | Heaviest rains in summer | Deciduous forests |
| | Subtropical | Coldest month between 18°C and -3°C , warmest month $>22^{\circ}\text{C}$ | Adequate throughout year | Coniferous and mixed coniferous-deciduous forests |
| | Marine | Coldest month between 18°C and -3°C , warmest month $>22^{\circ}\text{C}$ | Heaviest rain in winter | Coniferous forests |
| | Prairie | Variable | Adequate all year, except dry years | Tallgrass, grasslands, parklands |
| | Mediterranean | Coldest month between 18°C and -3°C , warmest month $>22^{\circ}\text{C}$ | Dry summers, rainy winters | Evergreen woodlands and shrubs |
| <i>Dry</i> | Steppe | Variable, winters cold | Rain $<50\text{cm/year}$ | Shortgrass, shrubs |
| | Desert | High summer temperatures, cold winters | Very dry in all seasons | Shrubs or sparse grasses |
| <i>Humid Tropical</i> | Savanna | Coldest month $>18^{\circ}\text{C}$, annual variation $<12^{\circ}\text{C}$ | Dry season with $<6\text{cm/year}$ | Open grassland, scattered trees |
| | Rainforest | Coldest month $>18^{\circ}\text{C}$, annual variation $<3^{\circ}\text{C}$ | Heavy rain, minimum 6cm/month | Dense forest, heavy undergrowth |

As all ecosystems operate within the context of larger systems, so our understanding of these large ecoregions allows us to better understand what is going on at a smaller local level. This knowledge provides a better foundation for ecological management of resources, such as plants, land, and water.



Ecological regions of North America: note that the province of Manitoba is quite diverse, and it includes the Great Plains (Prairie, Humid Temperate), Northern Forests (Warm Continental, Humid Temperate), Hudson Plain (Subarctic, Polar), Taiga (Subarctic, Polar), and Tundra (Subarctic, Polar).

PLANT COMMUNITIES

Ecosystems generally have well-defined plant associations. This plant association will be dependent on a variety of factors, including soil types, meteorology (rocks), and a mixture of fauna (animals). The plant association may change by seasons in temperate and boreal areas, although much of the appearance of difference will be due to dormant or leafless plants in the winter season.


STRATIFICATION AND STRUCTURE

In each ecosystem, a well-defined plant association is present, and often characterized by multiple layers. In the case of some ecosystems, such as grasslands, and other treeless habitats, some of these layers may be missing.

Vertical stratification

The structural features of plant communities, especially forests, are built on vertical stratification created by the amount of space taken up by specific vegetative groups. For example, in forests the amount of space occupied by trunks, branches, twigs, and leaves at different levels changes the amount of light spread throughout the various levels. These differences control the growth of plants adapted to each level. All together, these differences influence the plants and animals found at each level.

In general, four different strata are recognized. Not all layers are found in all locations. The **canopy** refers to the highest layer of vegetation in a forest, made comprised of the crowns of its tallest trees. Individual trees growing above the general layer of the canopy may form an **overstory**. The **understory** refers to those trees above the shrub layer and below the canopy. Light intensity starts to dim within this layer. The **shrub layer** is the layer of vegetation within a habitat with heights of between one and a half to about 10 metres. Young trees are also




part of this layer. This layer only receives light filtered by the canopy. Semi-shade or shade-loving plants that would not tolerate bright sunlight prefer this layer. At the edge of a forest, the shrub layer acts as a windbreak close to the trees and protects the soil from drying out. The final layer, the **floor** or **herbaceous layer** of a plant community, is composed non-woody vegetation, or ground cover, growing in the forest with heights of up to about one and a half metres. The herbaceous layer consists of various herbaceous plants, grasses, dwarf shrubs, and young shrubs. In deciduous forests, early flowering plants appear first before the canopy fills out. Thereafter, the amount of light available to plants is significantly reduced and only those that are suited to such conditions can thrive. In an oak forest, only about 6% of the mid-day sunlight will reach the forest floor. By contrast, grasslands consist of moss and herbaceous layers.

The floor of a plant community is often covered by a layer of dead animal and plant material. Dead trees, in the form of large standing dead trees, snags, or downed trunks and limbs also make up an important part of this layer. Through the work of microorganisms, bacteria, fungi, and algae, the organic material in this layer, from dead trees and other organic materials, nutrients are returned to the system. Further, this layer provides important habitat for animals that call the forest home. Under the floor of a plant community, the root layer (**rhizosphere**) are often also recognized. It consists of the plants' roots and related elements such as rhizomes, bulbs and tubers.

Slopes and Elevation

The plant community of an area can be strongly impacted by slopes and elevation changes within the landscape. The amount of sunlight and precipitation an area receives can change depending on the elevation,

consequently changing the composition of its plant and animal communities. As you can see in the figure below, the south-facing slope has plants reflecting an increased solar radiation and decreased precipitation. In contrast, the north-facing slope is composed of plants that need more water and precipitation.



HABITAT TYPES

Regions of the earth have also been grouped based on the types of vegetation present, known as vegetation regions, or habitat types. The four major types are grasslands, tundra, desert, and forests. Shrublands are a fifth area, that are often just examined as the transition between forests and other regions yet have their own unique vegetative structure.

Grasslands




© Phil Schermeister/National Geographic

Grasslands are the largest of the four major vegetational formations. They represent 24% of the Earth's vegetation. Native grasslands are found in regions with periodic drought, moderate amounts of rainfall, and the accumulation of organic matter. Grasslands are dominated by grasses. Grasses are distinctive because their stems produce narrow leaves that grow from their bases. This specific type of growth allows these plants to survive grazing by animals or be mowed. All grasslands have a climate characterized by high rates of evaporation, periodic severe droughts, and rolling-to-flat terrain. Forb species, including legumes, as well as an assortment of herbaceous plants (e.g., dewberry and goldenrod) are

often found in association with grasses in these areas. In prairies, such as in Manitoba, legumes and asters are important components to these areas.

Grasses are classified as either sod-formers or bunchgrasses. Sod-forming grasses grow a solid mat of grass over the ground. Bunchgrasses grow in bunches (see figure, with bunchgrass on left and sod-form on right). The space between bunchgrasses often can be filled by other plants including herbs. Little bluestem is a typical bunchgrass. Western wheatgrass is an example of a sod-forming grass, which is spread by underground stems. Some grasses can form either sod or bunch types depending on their environment. For example, Big bluestem will develop into a sod-form grass when the soil is rich and moist but form a bunchgrass when the soil is dry.



Growth forms and root penetration of a bunchgrass (left) versus a sod grass (right).
Modified from Smith and Smith 2001

Many of these natural grasslands have been converted into agriculture cropland. Grains such as corn, wheat, oats, and barley are often grown in these areas. Grasslands are also often used to produce hay or serve as pastureland for cattle or sheep.

Many types of grassland exist, with North American grasslands often being grouped into three major types, tallgrass, mixed-grass, and shortgrass prairie.

Cultivated and successional grasslands – Grasslands found in normally forested regions that are either cultivated (for agriculture) or successional. In North America in highly developed areas, cultivated grasslands are very common.



© Nicholas A. Tonelli

Tallgrass Prairie – Big bluestem is the dominant grass of the tall grass prairie, with little bluestem, porcupine needlegrass, and prairie dropseed grasses. A diversity of forbes are also found in this type of grassland. A large part of this grassland is interspersed with trees and shrubs, especially along streams and lower slopes of hills. Unfortunately, most of this prairie has been converted into cornfields and other cropland. It has been estimated that in Manitoba only 0.5% of the original tall grass prairie remains. Manitobans can visit some of the last remnants of this habitat type at the Manitoba Tall Grass Prairie Preserve.



Mixed-grass Prairie – Mid-height grasses occupy the lowlands and shortgrass occupy the higher elevations within the mixed-grass prairie. The amount of precipitation varies widely from year to year, which is reflected in the mix of grasses, forbes, and herbs.



© Anne Stine

Shortgrass Steppe – Shortgrass steppe/prairie contains shallow-rotted short grasses that reflect the reduced moisture found in this area. Sod-forming blue grama and buffalo grass dominate this landscape. Much of the short-grass steppe has been destroyed by overgrazing and wheat agriculture. The lack of moisture in this area meant the area could not support wheat agriculture for long and so

drought, lack of tight sod cover, and winds turned areas of the short-grass prairies into the Dust Bowl in the 1930's. Recovery from this period has taken decades.



© Llano Estacado

Manitoba's grasslands are part of the 'temperate grassland' zone that extends across the middle part of the United States and the southern parts of the prairie provinces. Temperate grasslands are the most endangered terrestrial (land-based) ecosystem on earth.

Shrubland



Shrublands are very diverse and range from Mediterranean types in arid and semiarid regions to heathlands of cool-to-temperate climates and successional shrublands. The moors of Scotland and the macchia of South America are two famous examples of shrublands. Shrubs and shrublands are difficult to characterize. Shrubs are often defined as a plant with wood stems, no central trunk, and a height of up to 4.5-6 m. However, size is not good characterizing measure and many areas called shrublands contain varied vegetation.

Shrubs are found to dominate areas experiencing low water, soils lacking in nutrients, cold winters, short growing seasons, and wind. Shrubs are able to take advantage of these area as they have less energetic and nutrient investment than trees, extensive root systems (take advantage of moisture deep in the soil), and many stems to influence the capture of water. Some species of shrubs may inhibit the growth of their competitors (such as herbs) by secreting substances toxic to other plants.

Deserts



Spruce Woods

© Solo Outdoors

Deserts occur where the annual rainfall ranges between 7-40 cm, it is possible that the rate evaporation will be higher than precipitation, and there is low plant growth. Woody-stemmed and soft-brittle-stemmed shrubs are characteristic desert plants. Yucca, cacti, and small trees all can also be found growing in the desert. Plants in this area have evolved to deal with the scarcity of water (xerophyte plants – see *Plant Adaptations to Extreme Environments*). Plants in this area either avoid droughts (drought evasion) or are drought

resistant (xerophytes). **Drought-evading plants** can be annuals (regrow each year) or perennial (persist over years). Often annuals in the desert persist as seeds during the drought and are ready to sprout, flower, and produce seeds when conditions change, and moisture and temperature are good. Drought-evading perennials have bulbs that send up growth during rainy periods, but they can bloom at different times. **Drought-resistant plants** all are perennials with different adaptations to this environment. Some shed leaves during the dry season while others maintain evergreen leaves throughout the year and have a taproot that reaches the water table. A further group that is often seen as symbolic of deserts is the succulents that have large internal water reserves to carry them through droughts.

Tundra



Tundras are areas that are cold due to high latitude or altitude. These areas are characterized by low temperatures, short growing seasons, and low availability of nutrients. The word tundra comes from the Finnish *tunturia* meaning “a treeless plain”. Arctic tundra can be found encircling the northern pole of Earth, whereas alpine tundra can occur at the peaks of tall mountains. In the Antarctic, a well-developed tundra is lacking due to the small amount of land area and deep ice sheets.

Tundra is land with many lakes connected by streams, and occasional rivers. The vegetation in this zone ranges from tall shrubs (2-5 m high) to dwarf shrub heath (5-20 cm high),

grasses, and mosses. In low lying areas extensive bogs can persist. In higher areas exposed to wind, vegetation is scarce and scattered, and the ground often is rock-covered and bare.


Permafrost, or a permanent frozen layer in the ground, as well as cycles of freezing and thawing shape the arctic tundra ecosystem. The depth of thaw may vary from a few centimetres to half a meter, which makes parts of the ground impenetrable for both water and roots. This creates shallow lakes and bogs. It also limits plant growth. However, the reservoir of water on top of permafrost also enables plants to be found in the driest parts of the arctic.

The vegetation of the tundra tends to be small, grow slowly, and most plants are from a few species' groups. Most plants in this area are perennial and grow vegetatively instead of using seeds. South of the arctic tundra and below the alpine tundra is a transitional zone between the tundra and the forest. The tree line marks the edge between these two habitats. Within this zone, **krummholz** or “crooked wood” trees are found. These trees are stunted and misshapen by the force of the wind, cold, and winter desiccation.



Forests

Forests represent the most widespread and diverse types of vegetation in the world. Distinct bands of forests grow around the northern hemisphere. Moving southward from the tundra, in order, coniferous, temperate deciduous, and finally around the equator, tropical forests are found. In the southern hemisphere, tropical forests are found more extensively.



Coniferous Forests

Coniferous trees are cone-bearing gymnosperms, from the group Pinophyta. They have dark green, needlelike or scalelike leaves. Other than a few exceptions, conifers are evergreen meaning they retain their foliage all year. Having foliage year-round allows the trees to photosynthesize throughout the year. Conifers are some of the most important source of lumber and paper pulp. A variety of coniferous forests can be found around the world.

Taiga and Boreal Forest: The taiga and boreal forest form a belt across the northern hemisphere. This forest is found within the polar domain. Severe winter is the dominant season and only a small amount of precipitation comes within the warm summer months.

However, due to the reduced evaporation due to cold, this type of forest can remain moist year-round. The evergreen coniferous trees found within this forest have adapted to these winters and respond quickly to the summer season.

Moss, lichens, and low shrubs are also important plants found within these forests. The exact composition of this type of forest varies throughout its extent. In Canada, these forests are dominated by four groups of conifers, Spruce (*Picea*), Firs (*Abies*), Pine (*Pinus*), and Larches (*Larix*), and two groups of deciduous (shed leaves annually) trees, Poplar, Aspen, and Cottonwood (*Populus*), and Birch (*Betula*). Jack pines and Black Spruce trees dominate in these areas.



Temperate Needle-Leaf Rain Forest: Found in the west coast of North America, within the humid temperate domain, these areas receive superabundant rainfall, high humidity, and warm temperatures creating unique forests. These areas are full of mosses, epiphytes and ferns, along with conifers that have adapted to these wet mild winters and nutrient poor soils. Hemlock, Pacific silver-fir, Douglas fir, dominate these forests. Coast redwood can be found in the southern areas of this forest. This type of rain forest forms the densest coniferous forests and contains some of the world's largest and tallest trees.



Coniferous Woodlands: Dry climates found in western North America, create the conditions for piñon-juniper woodlands. These ecosystems are part of the Temperate Desert Division.

These forests have open growth small trees in association with an understory of grass and shrubs. Coniferous woodlands have been impacted by livestock grazing, fuel harvest, and fire.




Temperate Broadleaf Forests



Despite their name, temperate forests face daily and seasonal temperature fluctuations, that shape this habitat. Deciduous trees drop their leaves annually, and so deciduous forests are leafless during the winter. Northern deciduous forests remain leafless most of the year. Yet, despite all this, temperate forests are highly productive, supporting high levels of plant and animal life.


Temperate Deciduous Forests: The temperate deciduous forests can be found worldwide, and once covered vast areas of Europe, China, South America, and eastern North America. Large parts of these forests have been cleared for agriculture and human development. Oak, beech, Ash, Birch, and Elms are the trees that dominate this type of forest. As these forests are highly productive, they are also home to a variety of other plants. Lichen, moss, ferns, wildflowers and other small plants can be found on the forest floor. A variety of plants fill in the shrub layer and trees make up the canopy.



Temperate Woodlands: The open oak and oak-pine temperate woodlands provide transition from coniferous forests to other habitats, including prairies, steeps, and deserts. Found in the southwestern United States and Mexico, this transitional woodland is found in dry areas. This type of forest is dominated by oaks, especially the Emory oak, but junipers and pines may be present as well.



Temperate Broadleaf Evergreen Forests: Found within several subtropical areas of the world, temperate broadleaf evergreen forests contain live oak, magnolias, and redbay. These forests are also characterized by many ferns and palms in the shrub layer of the forest, and epiphytes, lichen, and Spanish moss found throughout.



Tropical Forests



Tropical forests are found in the humid tropical domain, in the rain forest division. These forests experience a steady year-round temperature of about 23°C and a wide variety in rainfall. The variety in climatic conditions are reflected in the diversity of vegetation patterns found in these forests. The diversity found in these forests is unequalled anywhere else in the world.

COMPETITION

Plants must compete with each other, and other living things, for limited habitat resources, including space, food, nutrients, water, and light. Competition is the negative effects on plant growth or fitness caused by the presence of neighbours, usually by reducing the availability of resources. This competitive interaction occurs when a specific resource is in a short supply compared to the number of seeking the same resource.

Interspecific competition occurs when an individual or community is competing with another species, e.g., grass is competing with trees. **Intraspecific competition** occurs when an individual is competing with members of the same species, e.g., jack pine tree with another jack pine tree. Competition is an important factor that limits population growth as well as influences what species of plants and how many of them may be found in the same area.


Competition can be an important factor controlling plant communities, along with resources, disturbance, herbivory, and mutualisms. The effects of competition are widespread and easily observed in mixtures of crops and managed forests, which is why weeding and thinning are practiced. Competition is also widespread in native habitats, from deserts to wetlands, and is known to have important—indeed crucial—effects upon recruitment, growth, and reproduction.

SUCCESSION OF PLANT COMMUNITIES

Habitat, the complex association of soil, water and plants is in itself dynamic and ever changing. These changes can be subtle or dramatic. A forest fire causes a dramatic habitat change. The coniferous forest, cool and shady, disappears. Eventually, on the blackened, but now sunlit ground, grasses and other plants appear. Each type of plant appears, grows, matures, and disappears to be replaced by others, which also go through their stages and are replaced by still other varieties. This series of changes taking place is not random or haphazard but a predictable, sequential chain of events called succession. **Succession** is the orderly replacement of one biotic community with another. With each successional stage, be it subtle or dramatic, habitat is changed. With changes in habitat come changes in the forms of wildlife using that particular habitat.

In a recently exposed environment, such as an area that has recently undergone a major disturbance, like, severe fire, industrial development, or a volcanic, lichens and moss can serve as a pioneer plants moving into this area, followed by grasses and flower plants over time. This is slowly replaced by a low ground cover of grass and flowering plants. **Pioneer species** are hardy species that are the first to colonize an area, beginning a chain of ecological succession that ultimately leads to a more biodiverse steady-state ecosystem. Over the next few years – shrubs, bushes, or trees (e.g., willows, aspens and coniferous trees) each in turn, make their appearance. Finally, the plant community is once again as it was, composed with a **climax community**. This final or climax stage will remain until it experiences a disturbance, such as a fire, storms, or logging, and the successional cycle is triggered once again. Each species of wildlife has unique habitat requirements. Therefore, changes in habitat will change the kinds of wildlife associated with it.

Below is a simple example of succession:



COMMUNITY STABILITY AND EQUILIBRIUM

Succession can reach a climax in some communities, which produces a stable community that will be dominated by a small number of prominent species. In contrast, areas experiencing continual small-scale disturbances (e.g., fires, storms, etc.) will produce communities that are more diverse, and any species may become dominant for a time. Communities undergoing these fluctuations highlight the effects of unpredictable disturbances on the development of the community composition of an ecosystem. In some tropical forests where hundreds of tree species may be present in a small area, the death of a tree can shift the species composition within the same area. Other plants that previously could not live in this area due to a lack of resources, such as sun, nutrients, and water, can suddenly grow in this area.

Communities that are diverse, or have a large number of species, are considered healthy. If a community that is diverse undergoes a disturbance, such as fire, flooding, or storms, they

are faster at recovering than a community that has few species. Additionally, undisturbed communities can deal with the introduction of invasive species better.

Disturbance-dependent ecosystems

Disturbance is a natural part of many different ecosystems. Disturbances are often defined as “discrete events that disrupt the structure of a community or population and change resource availability.” Disturbances can be natural (not due to humans) and are an important part of how an ecosystem functions. Disturbances are important for nutrient cycling and the structure (how species are arranged) of an ecosystem. The frequency of the disturbance, how big the disturbance is, the intensity of the disturbance, and its severity all impact the role it plays in changing an ecosystem.

Non-natural disturbances can be caused by human development and actions. Humans have been modifying ecosystems for millennia, both directly (e.g., livestock grazing, fire suppression) or indirectly (e.g., landscape fragmentation, introduction of invasive species) or introduce new types of disturbances like pollution.


Minor disturbances can include localized wind storms, droughts, floods, small fires, and disease outbreaks due to pathogens. Major disturbances can include large-scale wind events, volcanic eruptions, hurricanes, intense forest fires, epidemics, weather events, pollution, and land use conversion (e.g., turning an area into agriculture or housing development).

Fires – Fires are natural disturbances that help shape the plant and animal communities throughout the world. Some ecosystems, like grasslands and some forests, are subjected to fires over centuries and we consider them to be **fire-dependent**. These communities require fires to restore and maintain their ecological integrity.

Fires produce a mosaic of plant communities with different ages and species composition within a landscape. Fires burn with a variety of intensities depending on differences in terrain, wind, and other factors. Typically, areas where the fire has completely consumed the landscape, are scattered within patches of lightly burned and unburned areas. The fire increases the diversity of structure (arrangement of species) and species diversity over time.

Nutrients can also be released during a fire. The fire helps release nutrients bound in litter and woody debris on the ground. Fire will also reduce woody fuels to ash and consume the organic layer of soil. Nutrients are lost to the atmosphere as smoke, but many more nutrients are added to the soil. This “flush” of nutrients is available to plants that are re-establishing themselves in burnt areas.

In the boreal forest, after a fire the burned sites will begin to regenerate with the establishment of pioneer species, such as white birch, jack pine, and lodgepole pine. These three species require full sunlight and are adapted to landscapes that experience regular fires. Jack pines and lodgepole pines have a type of cone known as **serotinous** which is protected by a waxy coating and requires heat to release their seeds. Fires produce the favourable conditions for these seeds to germinate and grow, and both pine species require fire to regenerate. Other species, such as balsam fir, white spruce, and white cedar have not adapted to fires and take a long time to recolonize areas that have experienced fire (sometimes as long as 150 years).



Fire regimes of different vegetation types in North America

© 2014 R.G. Bailey/Springer Science

The grasslands were created and maintained by the presence of frequent fires. Fires can remove encroaching trees and shrubs, as well as stimulating new growth of grass species, through the release of additional nitrogen and other nutrients. Ancient hunting peoples were also known to set regular fires to maintain and extend existing grasslands. Grassland plants, such as grasses, have long root portions that can survive the fires and sprout up again quickly. Some trees in these regions have thick bark to resist fire.


Herbivory – Plants are attacked and eaten by many different animal consumers. They form the base of the food web to which communities and ecosystems are assembled. Herbivories, in turn, can have a strong impact on plant growth, reproduction (directly as seed predators, indirectly by reducing the amount of plants), and survival. Herbivores can include zooplankton, snails, insects, waterfowl, fish, muskrats, moose, deer, etc. Large-scale studies have shown that herbivores can influence what plants occur within a community. For example, if a large mammal such as a group of deer move through a grassland, they may consume specific types of plants, giving more space for other plants to grow. In some environment's plants have adapted to extensive and consistent herbivory. In grasslands like the prairie, for example, grasses have extensive root systems to prevent grazing or herbivorous animals from pulling the roots out of the ground. They also grow from near their base, not their tip, and thus are not permanently damaged from grazing animals. These adaptations allow the grasses to regrow quickly after being consumed by animals such as Bison or cattle.



The difference grazing by a large herbivore (cattle) can make on plant communities

Flooding and Water Levels – Wetlands, such as marshes, bogs, and fens, need fluctuating water levels to function properly. These fluctuating water levels are necessary to maintain dynamic, diverse, and healthy wetlands. Cycles of high to low water levels help create the diversity in vegetation that is more resilient to other disturbances. High water levels help prevent upland woody plants from invading and increase the meadow marsh at higher elevations. Low water levels allow the seeds on the water bed to germinate, allowing the establishment of many emergent plant species. Low water levels also reduce the invasion of meadow marsh by moisture requiring cattails. Water-level change patterns are the driving


force determining the overall diversity and condition of wetland plant communities and the habitats they provide for invertebrates, amphibians, fish, birds, and mammals.



ECOTONES AND EDGE EFFECTS

The **ecotone** is a transitional area of vegetation between two different plant communities, like a forest and grassland. This area will have some of the characteristics of both habitats but could also have plant species not found in either overlapping community. **Edge effects** are the influence of the two bordering communities on each other. The ecotonal area often has a higher density of organisms of one species and a greater diversity (number) of species than either habitat alone. This area is crucial for some organisms for courtship, nesting, or foraging.

Ecotones can also be found where one body of water meets another (for example estuaries or lagoons), or where water meets land (for example wetlands). Both marine and freshwater ecotones have large plants that rise from roots attached to the submerged substrate.



PLANTS AND THEIR ENVIRONMENT

Plants have adaptations to help them survive (live and grow) in different areas. Adaptations are special features that allow a plant or animal to live in a particular place or habitat. These adaptations might make it very difficult for the plant to survive in a different place. This explains why certain plants are found in one area, but not in another. For example, you wouldn't see a cactus living in the Arctic. Nor would you see lots of really tall trees living in grasslands.

WINTER ADAPTATIONS

Plants have evolved a variety of adaptations to surviving the winter. Unlike many animals (and humans!), plants cannot leave, hibernate, or otherwise escape the winter conditions. As they are rooted, plants must stay where they are, and so they have evolved strategies to survive the conditions around them.

Shedding Leaves

Deciduous plants survive the winter and its lack of water by shedding their leaves, as water tends to get evaporated into the air from the leaves. During the fall, leading up to the winter, most deciduous trees drop their leaves and become dormant. See the figure for more details on this process.

Evergreen trees (like



Why do leaves change color in the Autumn?

Most of the spectacular colors of autumn have actually been in the leaves all summer, however they were "covered up" by the dominant green of the chlorophyll. As weather cools, and shorter days settle in, the chlorophyll begins to break down, revealing new and varied color pigments. The brightest colors are seen when late summer is dry, and autumn has bright sunny days and cool nights.



GREEN - Chlorophyl

Chlorophyll is responsible for helping trees and plants turn sunlight into food. For most months, it is the dominant color seen in most leaves until it fades away. As many trees shut down their food production, they turn to stored sugars to survive the winter.




RED - Anthocyanin

Unlike other leaf colors that always exist in the leaf, anthocyanins are produced as the chlorophyll is broken down. The anthocyanins are often seen in leaves named for their autumn splash of red including Red Maples, Scarlet Oaks, and Red Sumacs.



ORANGE - Carotene

Sugar Maples may be one of the best examples of carotene in action. Their bright signature orange fills many hills and country roads throughout the northern US. Sassafras leaves also turn a slightly more muted orange. As its name implies, Carotenes are also the chemical responsible for giving carrots their unique coloring.



YELLOW - Xanthophyll

Xanthophyll can be seen throughout the fall in trees including beeches, ashes, birches, aspens, and some oaks. It also contributes its bright yellow color to autumn squash and corn.

Please print and share! - facebook.com/sciencebob

coniferous trees) maintain their leaves and needles. These leaves and needles have a thick, waxy coating that reduces water loss.


Plant Growth and Shape

Areas that are cold year-round and receive frequent snow, such as the Arctic, have trees that grow close to the ground or grow in shapes that help them shed snow easily, such as branches that droop downward to allow excess snow to fall to the ground and prevent breakage. Plants in this region may also hold onto their leaves to help insulate them from the cold.

Plants within cold environments also go dormant during the winter period to reduce damage from temperatures and frost. However, if sharp frost events following warm spells or when snow bed plants become snow free may lead to some tissue loss by plants. This type of characteristic damage can be observed in wood products.

Trees in cold locations, such as the Taiga, are dominated by conifers. Coniferous trees are evergreen which allows the plant to photosynthesize as soon as the temperature rises to a minimum temperature. Further, the needles are dark in colour allowing more solar heat to be absorbed.

In the Tundra, which is cold year-round, plants have also developed the adaptation to be covered with a hair like growth that helps insulate them. Plants also grow in groups to protect each other from the wind and cold.



Freezing Resistance

In order to survive the cold, plants must have some form of resistance to freezing. Changes within a plant allow it to survive temperatures below the freezing temperature of water, although many of these processes require a slow acclimation period. For example, some coniferous trees also have a special valve in their cells that can seal off individual frozen cells to prevent a freezing chain reaction. Additionally, some trees in cold regions are known to be able to super-cool their xylem water (cool down to -40C).

WIND ADAPTATIONS

Stems and Leaves

Plants that have to undergo great stress due to constant windy conditions often have evolved to have soft stalks, so they can flex with the wind instead of fighting against it. For example, the grasses found in a grassland are extremely flexible and can bend in the wind.

The leaves of some species of trees deal with strong winds by adjusting their configuration (curling) in order to limit flutter and their exposure.



Seed Dispersal

Some plants take advantage of living in windy environments, using the wind to distribute their seeds throughout the environment.

Size and Clustering

Some plants, such as those found in the tundra, will cluster together to protect each other from the wind. Additionally, in this area plants will remain small to reduce the impacts of wind.

PLANT ADAPTATIONS TO EXTREME ENVIRONMENTS

Plants are adapted to the location to which they live. In nature, situations can change. Sometimes areas may become dryer, wetter, or sometimes taken over by a specific organism. These changes can be rapid, but often they have occurred over a long period of time. Plants and animals have mutations in their genetic code, which can be harmful, neutral, and sometimes beneficial when conditions change. In ecology, an **adaptation** is a change that is maintained by natural selection.

Changes in environmental conditions for plants could be caused by a variety of factors, like decreased availability of water (drought) or excess water within its habitat (flooding). Plants that have adaptations that are better suited to this changed environment may do better than those without any adaptations, and over long periods of time, the whole new population will inherit this change. In areas that experience these extreme conditions over millennia, plants have evolved extreme adaptations to deal with the extreme conditions.

ECOLOGICAL GROUPS OF PLANTS: HYDROPHYTE, XEROPHYTE, MESOPHYTE, HALOPHYTE

Plants often have adaptations, or changes, that help them survive under a range of conditions.

Hydrophyte – this group of plants are adapted to living in aquatic environments. These plants either remain fully submerged in water, like *Hydrilla*, or most of the plant remains under water, like lotus or water lilies.

Xerophyte – this group of plants are adapted to survive in an environment with little liquid water. Examples include cacti and succulants.

Mesophyte – this group of plants are adapted to a habitat with adequate water. These plants include most of the plants we see around us every day, like trees and shrubs.

Halophytes – this group of plants are adapted to a salty habitat. These plants are often found in saline sem-deserts, mangrove swamps, marshes, and sea shores. Examples include cordgrass, switchgrass, and saltbush.

HYDROPHYTES

- Underwater leaves and stems are flexible to move with water currents.
- Some plants have air spaces in their stems to help hold the plant up in the water.
- Submerged plants lack strong water transport system (in stems); instead water, nutrients, and dissolved gases are absorbed through the leaves directly from the water.
- Roots and root hairs reduced or absent; roots only needed for anchorage, not for absorption of nutrients and water

- Some plants have leaves that float atop the water, exposing themselves to the sunlight
- In floating plants chlorophyll is restricted to upper surface of leaves (part that the sunlight will hit) and the upper surface is waxy to repel water
- Some plants produce seeds that can float

Waterlilies leaves, as you can see in the cross-section, have a tough network of bundled hollow cells that hold air that helps the stems float.





Figure 25-21
Biology of Plants, Seventh Edition
© 2005 W.H. Freeman and Company

XEROPHYTES

A xerophyte is a species of plant that has adaptations to survive in an environment with little liquid water, such as a desert or an ice- or snow-covered region in the Alps or the Arctic. Popular examples of xerophytes are cacti and pineapple plants.

- Some plants, called succulents, store water in their stems or leaves
- Some plants have no leaves or small seasonal leaves that only grow after it rains. The lack of leaves helps reduce water loss during photosynthesis. Leafless plants conduct photosynthesis in their green stems.
- Long root systems spread out wide or go deep into the ground to absorb water;
- Some plants have a short life cycle, germinating in response to rain, growing, flowering, and dying within one year. These plants can evade drought.
- Leaves with hair like structures help shade the plant, reducing water loss. Other plants have leaves that turn throughout the day to expose a minimum surface area to the heat.
- Spines to discourage animals from eating plants for water
- Waxy coating on stems and leaves help reduce water loss.
- Flowers that open at night lure pollinators who are more likely to be active during the cooler night.
- Slower growing requires less energy. The plants don't have to make as much food and therefore do not lose as much water.



Mechanisms of drought resistance

© from Brunner et al. 2015, Frontiers in Plant Science

COMMON ENVIRONMENTAL PROBLEMS

Plants often experience similar environmental problems, especially with anthropogenic or human development.

Foliage burn

Foliage burn or leaf scorch can be caused by soil compaction, transplant shock, nutrient deficiencies, drought, salt toxicity, and herbicide injury. Foliage burn is defined as a browning of plant tissues, including the margins and tips, and a yellowing or darkening of veins which may lead to eventual wilting and release of the leaf. To reverse these symptoms and the damage, pruning sprouts and affected areas, avoiding frequent light watering periods and watering heavily to promote healthy root development, and avoiding over-fertilization can be used.



Pollution

Pollution is a worldwide problem. It is wide-spread, and it can come in many forms, including water, air, and land (soil). Sources can include industry, commercial development, and transportation. Pollution can affect humans and animals, but also plants. Damage can include leaf damage, slower growth, root damage, and an inability to properly photosynthesize.

Air pollution – air pollution can come from a variety of sources, including a factory, car exhaust, or off gassing from the production of plastic. The impacts of air pollution on plants are wide spread. Damage in plants can be observed in a variety of ways including lesions, stunted plant growth, or the changing of colour (aka yellowing of leaves, reddening, bronzing). The holes in the ozone can also be very damaging for plants. Holes in the upper atmosphere let excess UV light to pass into the atmosphere which leads to plant damage. In the lower atmosphere ozone damages plants by preventing photosynthesis and blocking stomata, which restricts respiration and stunts plant growth.

Soil pollution – This type of pollution can come from improper waste disposal, oil spills, landfills, pesticides, or the illegal dumping of toxic substances. The addition of chemicals into the soil can strip it of nutrients as well as fill the soil with chemicals or metals that may damage plant cells and limit growth if they are taken up. If the chemicals are toxic they can also change plant growth and reduce crop yields.


Water pollution – Water pollution may be present due to sewage leaking, industrial spills, biological contamination, or farm runoff. Contamination of water will have many negative effects on plants including excess nutrients (excess plant growth), fluctuations in acidity that can damage or kill a plant, and finally, as with soil pollution, the uptake of chemicals.

Lack of water

Vascular plants have a system that enables water and nutrients to be taken up from the environment through their complex root system. The water needs of plants do vary widely. The species and age of the plant, the soil where it is rooted, and its sun exposure all are important factors. Some plants also have adaptations to exist in low water conditions. However, most plants need a continual flow or presence of water and nutrients to ensure the vascular system remains firm and the plant continues to be healthy. The lack of water or severe reduction in this resource can lead to many problems. For example, the lack of water can cause wilting of stems and leaves. Plants may also close down areas of the vascular system to reduce water loss, which will lead to leaf, flower, and fruit loss. Continued water stress can also make a plant more vulnerable to insect pests and disease, as well as less effective competitors. Although most plants have adapted to go through short periods of reduced water availability, sustained periods of drought can lead to the death of plants not adapted to low water conditions.

How tree roots respond to drought

The response of tree roots to drought involves a variety of strategies. Plants have evolved strategies that enable plants to avoid low water potentials and adaptations that enable them to tolerate low water conditions. Tree roots can adjust their biomass, change their anatomy (like increasing water pores), and acclimatize to the conditions. Mutualistic relationships between the tree and mycorrhiza can help a tree survive drought conditions.



Shade

Plants need light to photosynthesize and grow. Shade, naturally from other plants (like trees), or by buildings or other anthropogenic structures can have a negative effect on plant growth. Yet, plants may have a variety of adaptations to survive and flourish when experiencing shade. Plants that thrive in the shade, or shade-tolerant plants have adapted to conserve energy and optimize their use of any energy they have. Shade-tolerant plants have more chlorophyll in their leaves, lens-shaped cells that can focus incoming light, invest more in herbivore defense (producing toxins), and flowers and fruit that are almost invisible as they match the surrounding plant.

Humidity and temperature

The overabundance of water in the air can cause problems such as promoting the growth of bacteria and fungi which could be harmful to plants. Hot temperatures can also increase evapotranspiration of water and stress plants.

In ecosystems where the temperature is hot, and it rains a lot, like the tropical rainforest, plants have been able to adapt to this high humidity and temperature environment.

- Drip tips and waxy surfaces allow water to run off, to discourage growth of bacteria and fungi
- Buttresses and prop and stilt roots help hold up plants in the shallow soil
- Smooth bark and smooth or waxy flowers speed the run off of water
- Plants have shallow roots to help capture nutrients from the top level of soil.
- Many bromeliads are epiphytes (plants that live on other plants); instead of collecting water with roots they collect rainwater into a central reservoir from which they absorb the water through hairs on their leaves

ENVIRONMENTAL CONSERVATION

PESTICIDES AND FERTILIZERS

Pesticides and fertilizers can often be used to help plants grow.

Pesticides are used to kill off insects and other pests that might consume or negatively impact plants. Historically they have been used in agriculture to increase crop yield and to

improve pasture and grassland conditions. However, many pesticides can have negative impacts on the environment. Some may kill animals that are detritivores and an important part of the nutrient cycling within an area. If these animals are absent, nutrient cycling may slow down substantially, starving plants of the necessities of life. Further, as many pesticides are not very specific, they may also impact insects and other invertebrates that are important pollinators. Without these pollinators, many plants will be unable to reproduce.

Pesticides may also pollute the air, water, and soil. As a result, plant and animal life may become sick, malformed, or even die.

Fertilizers are materials that are applied to soils or plant tissues to supply one or more essential nutrients to assist with plant growth. By adding additional nitrogen and phosphorus to the soil, plants can thrive, as well as increase growth and reproduction.

However, there can be too many nutrients added within a system. Excess nutrients can become a type of pollution and wash off the soil into aquatic environments leading to eutrophication and other problems. Phosphorus from fertilizer can cause algae to accumulate in lakes and ponds, leading to decreased oxygen and the death of aquatic animals such as fish.

INVASIVE SPECIES

An **invasive species** is an exotic (originating from another region of the world) species whose introduction causes or is likely to cause economic harm, environmental harm, and/or harm to **native species** (including human) health. Species include plants, seeds, eggs, spores, other propagules, and animals (e.g., mammals, reptiles, amphibians, fish, insects and other invertebrates). This expansion is often due to human activities. Invasive species are more commonplace than one might think. Kentucky bluegrass, periwinkle, lily of the valley, and dandelion are all common plant species found in our lawns and gardens but are invasive species to this region. The domestic cat is thought to have originated in Africa. Some species have moved within the country into areas they have been previously absent. For example, the house finch, native to several western provinces, is now found in a number of eastern provinces.

Although all invasive species are **non-native species**, not all non-native species are invasive. Non-native species are only considered invasive if they have harmful ecological, environmental, or economic affects. All ecosystems are at risk from the harmful effects of invasive species. The adverse effects of invasive species do vary widely, from the extirpation or extinction of native species to small long-term effects on ecosystem function.

Invasive species' grow and reproduce rapidly, causing major disturbance to the areas in which they are present. These species can threaten an area's biodiversity by overwhelming native species, damaging habitat, disrupting food sources, and introducing parasites and disease. Most invasive species have little to no population control mechanisms in place to help control their population levels in the area of introduction and therefore often increase in numbers rapidly. Once invasive species are established in a region they can be difficult, or impossible, to control and remove.

Invasive species often share characteristics that make them successful in their new region. Invasive species characteristics include:

Few natural enemies

Many invasive species do not have any natural enemies (e.g., predators, competitors, parasites, and pathogens) in the area they invade. A lack of predators and pathogens may allow the invasive species population to spiral out of control.

High reproductive rates

Invasive species frequently have rapid growth, very short life cycles, prolific young production (e.g., prolific seed production), and seed dormancy (in plants).

High survival

Invasive species often can tolerate a wide range of environmental conditions. Invasives often can use a variety of pollinators (e.g., insects (such as bees, wasps, butterflies, etc.) and birds) to complete their life cycle.

Good dispersal

Most invasive species can very effectively distribute themselves into new environments. A lack of natural barriers, predators, and intraspecific competition may allow them to spread quickly throughout the new region.

Aggressive competitors

Most invasive species are superior competitors to native species. They may be more effective at obtaining resources like food, water, and/or space, or be better specialized at obtaining one specific set of resources.

A combination of these characteristics allows invasive species to outcompete native species in a region and become established.

Invasive species come into Canada by any means of transport that moves them farther than they could move on their own. Sometimes they are brought in on purpose, but often they

arrive unintentionally. Seafaring European explorers and settlers were the first to introduce new species to Canada. They brought cattle, goats, and other domestic animals, along with familiar crops like wheat, when they came by ship to explore and settle the New World. Without meaning to, they also introduced unwanted organisms—pests, like the Norway rat, and viruses, like deadly influenza and smallpox.

Many invasive species are transported to an area by accident. Accidental arrivals are rarely discovered until they have established themselves and have spread beyond their point of entry. For example, seeds of various plants can accidentally be transported from one location to another.

When an invasive species enters an ecosystem, it can have an impact on the species that are present, on important habitats, or even on the ecosystem itself. Concern arises when an invasive species changes the system for the worse, by either reducing or eliminating populations of native species, or by otherwise changing the way the ecosystem works. These changes have made the invasion of alien species a major global problem. If organisms were not able to move beyond their normal ranges, each part of the world would have a unique array of plants, animals, and microorganisms. However, as species move from one area of the world to another, sometimes squeezing out the competition, different places in the world become more alike in their biology—a process called **biological homogenization**.

The accidental introduction of the invasive Asian chestnut blight fungus almost completely eliminated American chestnut from 180 million acres of eastern United States forests. The introduction, through the nursery trade, caused an entire transformation of the Eastern deciduous forest ecosystem. The loss of the American chestnut was a disaster for many animals that were highly adapted to living in forests that were dominated by this tree species. Eventually, ten moth species that had relied on chestnut trees became extinct.

Garlic mustard is an invasive forest plant species native to Europe, introduced to North America in the 1800s to use as an edible herb. Garlic mustard is an allelopathic ground cover plant, which allows it to out-compete native ground cover plants and reducing overall ground cover diversity in forests. Its ability to reproduce rapidly expands its distributions in newly settled environments. While there



American Chestnut

© American Chestnut Foundation



Garlic Mustard

are methods of control including chemical and manual, timing of control is essential in trying to decrease the potential spread of garlic mustard.

European buckthorn was introduced from Eurasia in the 1800's to be used for urban landscaping and in parks. The invasive buckthorn has come to dominate mid-level canopies in many disturbed urban forests throughout North America. It has a high fecundity and a prolific growth rate. The success of buckthorn has been accelerated by a lack of natural controls and additional invasive species.



European buckthorn

Leafy Spurge - this plant is probably the most serious invasive in native grasslands in Manitoba. Leafy spurge (*Euphorbia esula* L.) is a deep-rooted perennial plant, which can reproduce by seed and underground creeping rootstocks. The plant stands approximately 50-60 cm in height, has yellowish-green flowers, contains milky white latex, and is usually found growing in patches. Infestations of leafy spurge generally occur in grasslands. The noxious plant often renders pasture lands useless for grazing as the milky latex causes detrimental effects to most grazing animals. However, sheep and goats appear to be unaffected and they will eat the plant. Nevertheless, losses in grazing land for cattle in Manitoba have been estimated at over half a million dollars per year. Leafy spurge is probably the most difficult invasive plant to control in Manitoba. As it has a well-developed root system, the plant is able to survive a number of different control methods (i.e. chemical and mechanical).



Leafy Spurge

New tool to predict which plants will become invasive

November 8, 2018 University of Vermont

<https://phys.org/news/2018-11-tool-invasive.html>

Around the world, over 13,000 plant species have embedded themselves in new environments—some of them integrate with the native plants, but others spread aggressively. Understanding why some plants become invasive, while others do not is critical to preserving the world's biodiversity. traits that are different from the native community and that plant height can be a competitive advantage.



New research from the University of Vermont provides insight to help predict which plants are likely to become invasive in a particular community. The results showed that non-native plants are more likely to become invasive when they possess biological

"Invasive species can have a devastating effect on our natural ecosystems and cause long-term environmental and economic problems," said Jane Molofsky, a professor in UVM's Department of Plant Biology and senior author of the study published November 6, 2018 in *Nature Communications*. "Our aim was to leverage big data and statistical techniques to evaluate this problem in a novel way by comparing traits of native and non-native plants across a range of plant communities."

Working with a team of international collaborators, Molofsky and colleagues at UVM explored differences in biological traits of 1,855 native and non-native plant species across six different habitat types in temperate Central Europe.

In each habitat type, the authors compared the traits of native and non-native plants. Of the non-native plants, they looked at differences in those that "naturalized," meaning they reproduce in nature without direct intervention by humans but did not aggressively spread, and invasive species, those that spread over long distances and often cause serious ecosystem damage.

Being taller promotes success

In almost all of the studied habitats, the findings showed non-invasive plants shared similar traits with the native plant community, such as plant height, leaf characteristics and average seed weight. In contrast, invasive species appeared to have similar but slightly different biological characteristics—they were similar enough to be present in the same habitats but just different enough to have unique characteristics that allowed them to flourish.

For instance, some invasive plants were taller on average compared with the native species. This phenomenon suggests the additional height of some invasive plants gives them better access to light and enables them to outcompete native plants and spread more aggressively.

The findings support a novel theory of invasion called the edge of trait space model that suggests non-native plants can co-exist with a native plant community when they share a set of specific biological traits but can invade when they have slightly different adaptations to local environmental conditions. Therefore, newly introduced species must be similar enough to thrive in a community of native species, but their differences may enhance their invasion success.

The results indicate that a single, easily measurable trait—plant height—can be a highly predictive factor in determining which plants may become invasive in a given environment. While the predictive traits may differ among different flora, the research suggests eradication efforts should focus on non-native plant species that differ from their native communities.

"We need new predictive tools to help inform policy and management decisions around conservation and biodiversity," said co-author Brian Beckage, a professor in the Department of Plant Biology and Department of Computer Science, and affiliate of the UVM Gund Institute for the Environment. "Our hope is that this model can be used as a screening tool to determine which plants have the highest probability of becoming invasive in the future."

More information: Jan Divíšek et al. Similarity of introduced plant species to native ones facilitates naturalization, but differences enhance invasion success, *Nature Communications* (2018). DOI: 10.1038/s41467-018-06995-4

Journal reference: *Nature Communications*

Provided by: University of Vermont

Invasive species can be added to a community either by natural range extensions or because of human activity. Humans have served as both unintentional and deliberate dispersal agents for millennia. In the last 200 to 500 years, the increase in human movement and trade has dramatically increased this dispersal. Human activities may include international, national, and regional trade and travel, horticulture, gardening and ornamentals, transportation and unity corridors, seed mixtures (re-vegetation, bird seed, wildflower), recreation, wildlife, livestock, humans, and pets (including the pet trade).

Cheat grass was introduced to North America in 1889 through shipments of grain seeds from Europe. Wooden packing material is often used to protect shipments of goods. These materials can often harbor invasive plant pathogens and insects. The Asian long horned beetle has been intercepted in wood packing materials in the USA and the UK.



Cheat Grass

© Robb Hannawacker/NPS

Time to Rethink the Lawn

Ecology and Field Biology
Smith and Smith 2001

It's any weekend day from April to October. America heads for the garage or garden shed to start the lawn mower. The tasks are well defined: mow, weed, fertilize, lime, aerate, and irrigate. The enemy is identified: crabgrass, dandelions, grubs, moles, ants, and Japanese beetles are among their ranks. The attack begins with herbicides and insecticides. It is an ongoing crusade, and the Holy Grail is the perfect lawn—the envy of the neighborhood.

The reason for America's obsession with the lawn is not clear, but theories abound. Some say it's rooted in early 19th-century British ideas of natural beauty. Evolutionary psychologists suggest that the love of lawns is genetically encoded—a molecular memory of the time when progenitors of the human species left their arboreal dwellings for Africa's savannas. Whatever the reason, the lawns of 58 million American households combine to cover 20 million acres of the United States—an area roughly the size of Pennsylvania. Nor is this obsession inexpensive. Americans spend an estimated \$30 billion per year on lawn care (enough to provide 1.5 million families with an income above poverty level). Despite the enormous economic expenditures, the real cost of creating and maintaining our lawns is perhaps best measured not in dollars but by the cumulative effects on environmental and human health.

Lawns are human-made ecosystems. To create a lawn, a bare piece of land is required. In the eastern regions of the United States, this means clearing forests. Most of the nitrogen and other essential nutrients in a forest are in the living vegetation and dead organic matter on the soil surface. When these are removed (by bulldozing or burning), chemical fertilizers must be applied in order for the topsoil to support a lawn. The nutrients in synthetic fertilizers are in an inorganic form that is readily available for uptake by grass plants. However, because these nutrients are not easily stored in the soil, they leach into ground or surface water. In contrast, nutrients in the organic matter that was previously on the surface of forest soil became available slowly through decomposition and nutrient mineralization (the transformation of nutrients from organic to inorganic form). Very little was lost from the ecosystem through leaching.

The problem of nutrients leaching from denuded forest soil is compounded when homeowners remove mowed grass clippings for aesthetic reasons. Removal of

clippings may result in a loss of up to 100 pounds of nitrogen per acre of lawn per year. Without the natural recycling of nutrients via decomposition of grass clippings, additional fertilizer needs to be added to maintain the lawn. Furthermore, to ensure maximum productivity, lawn owners often add more fertilizer than plants are capable of assimilating. This practice often leads to beautiful lawns, but it can create serious environmental problems.

Excess nutrients have a variety of negative effects on a lawn. Too much nitrogen can increase a grass plant's vulnerability to disease, reduce its ability to withstand extreme temperatures and drought, and discourage the activity of beneficial soil microorganisms. In addition, some synthetic fertilizers acidify the soil, reducing the uptake of magnesium, calcium, and potassium (elements important to biological and chemical processes) in plant and soil organisms.

The environmental impact of overfertilization is not limited to the lawns; it also affects ecosystems that communicate with lawns through the exchange of energy and matter. Nitrous oxide, a product of the breakdown of ammonia (a form of nitrogen commonly used as lawn fertilizer), is implicated in contributing to global warming. When leached from the soil into waterways, the nitrogen and phosphorus in fertilizers can lead to the eutrophication of aquatic ecosystems. This process begins with excessive growth of water plants and ends in a smelly body of water deprived of oxygen and the loss of many life forms. Chemical contamination of wells is a notable problem for humans, with nitrate from lawn fertilizers being the most common pollutant. Recent EPA surveys indicate that 1.2 percent of community water systems and 2.4 percent of rural domestic wells nationwide contain concentrations of nitrate that exceed public health standards. High concentrations of nitrate in drinking water may cause birth defects, cancer, nervous system impairments, and “blue baby syndrome,” a condition in which the oxygen content in the infant’s blood falls to dangerously low levels.

Further environmental damage occurs from lawn owners’ adversarial relationships with nature. In a lawn owner’s eyes, three groups of organisms threaten the lawn: animals (such as moles and insects), weeds, and fungi. They may bring disease or simply change the appearance of the lawn, disrupting the smooth, even carpet of green. Since the 1950s, many homeowners have waged war on these enemies with pesticides, specifically with rodenticides, insecticides, herbicides, and fungicides. . In contrast, naturally occurring ecosystems protect themselves against disease and insect outbreaks in many different ways. Some plants, such as milkweed, produce chemicals

in their leaves that make them unpalatable. Many insect populations are held in check by predators or diseases that are absent—due to pesticide use—in lawn ecosystems. Although pesticides may hold down populations of unwanted lawn pests for a time, they also result in the death of beneficial organisms—such as birds, earthworms, some insects, bacteria, and fungi—important to the health of the lawn. In addition, pesticides may persist in the environment for a long time with their lethal capabilities intact, even as they travel by wind, surface runoff, or seepage through the soil to wells and reservoirs used for public water supplies, wetlands, streams, rivers, and lakes, and even to marine environments.

Pesticide contamination of groundwater is less documented than fertilizer pollution but is of growing concern. Detectable levels of pesticides or their chemical breakdown products have been found in 10 percent of the wells in community water systems. Of the various types of pesticides, the most is known about insecticides. Many insecticides work by blocking communication between cells of the nervous system, but the newest ones, such as Bt (for *Bacillus thuringiensis*), prevent insects from maturing. The latter are considered to be nontoxic to vertebrates because they are naturally occurring insecticides that function by affecting the insect's growth hormones.

However, we do not know their long-term effects when used liberally, and we ought to keep in mind that many invertebrate hormones are similar or identical in molecular structure to vertebrate hormones. Furthermore, history has shown that when we use nature's products for our own purposes, rarely do we understand the full implications of our interference with the environment.

Left unattended, our lawns would quickly be invaded by a variety of native plant species, eventually giving way to the ecosystem that formerly occupied the site. To stave off the inexorable processes of succession, herbicides are applied to lawns. Only recently has it become apparent that herbicides, like fertilizers and insecticides, are also cycling throughout the environment. Because the data are both scarce and new, the public health implications of these findings are unclear. Nevertheless, herbicides are among the suspects for the suddenly high incidence of deformities observed in amphibians.

Only 5 percent of total yearly herbicide use in the United States is attributable to homeowners, but this 5 percent amounts to an astounding 33 million pounds. Herbicides come in a multitude of formulations. Some are selective, killing either


broad-leaved plants or grasses, but not both; others are nonselective, killing all vegetation. The most toxic herbicides cause death if ingested by damaging cellular components; others prevent the germination of seeds. Herbicides work by a variety of means including inhibiting cell division; electron transport systems; or the synthesis of lipids, chlorophyll, or key enzymes. Because traces of herbicide residues are being detected in unexpected and unwanted places, including our drinking water, we might be wise to exercise caution in their application until enough time has passed for us to understand their cycling in the environment and their potential toxicity.

Another problem with lawns is that they need watering. The United States is a nation where population increases have combined with increased per capita consumption of water to generate a water crisis. Since 1950, the rate of public water use (which excludes agricultural and self-supplied industrial use) grew at more than twice the rate of our population increase. Water tables are falling, and stream flow is decreasing in many river basins. For instance, water tables in the Dallas–Fort Worth area have fallen more than 400 feet in the last 30 years. Some states in the West are now battling over water rights among themselves and with Mexico, and water shortages are now also frequent on the typically much wetter East Coast. With lawn watering accounting for up to 60 percent of urban water use in the West and 30 percent in the East, it's not surprising that an increasing number of communities are restricting the use of water for lawns.

Grass grows from below ground in what is probably an evolutionary adaptation to grazing animals. We respond by using mechanized grazers—lawnmowers—to keep the grass at a uniformly short height. This, among other reasons, is why lawns are solidly linked to environmental issues that surround fossil fuel consumption, including smog, acid rain, oil spills, destruction of the ozone layer, and global warming. The most obvious use of fossil fuels for lawn management is in running the fleet of mechanized equipment, including mowers, aerators, leaf blowers, weed whackers, and edgers. A staggering 580 million gallons of fuel is consumed annually just by gasoline-powered mowers alone.

Fossil fuels are also employed to manufacture and transport inorganic fertilizers and pesticides. The principal nutrients in lawn fertilizers are nitrogen, phosphorus, and potassium. Natural gas is a reagent in the production of ammonia, the most common source of nitrogen in inorganic fertilizers. Furthermore, industrial nitrogen fixation requires a great deal of heat and pressure, which are supplied by energy from fossil fuels. Fossil fuels are also used to mine and refine potassium and phosphorus. The bag

of fertilizer you might buy in Virginia may have originated, in part, in Peru, Utah, and Saudi Arabia. Still more fossil fuel is required to power the ships, trains, trucks, and cars to move these products from their source to the lawn. The same is true for pesticides. Thus, calculation of the environmental costs of these products involves not only their direct chemical effects but also their hidden costs, such as consumption of fossil fuels.



Types of native grasses compared to Kentucky Blue Grass

When we think of managed ecosystems, we typically envision agricultural fields or forest plantations, yet the American lawn is the most expensive and management-intensive of all ecosystems. The endless nature of our lawn maintenance cycle is clearly evident: water and nutrients promote growth; herbicides lessen competition from weeds; rapid growth requires frequent mowing; pesticides inhibit not only grass predators but also organisms that decompose clippings; to achieve the desired appearance, clippings are removed; the removal of clippings requires fertilizers to replace nutrients, which promotes growth, etc. Changes are underway, with landscapers attempting to incorporate more native species of grasses and other herbaceous vegetation adapted to the local environmental conditions and requiring less maintenance. It is now becoming fashionable to create meadows and mixed-plant gardens that will attract wildlife in place of unbroken expanses of lawn (Figure 2). Research is ongoing to find nontoxic pesticides that do not easily cycle through the environment. Growing concerns about environmental pollution, energy conservation, and scarcity of water are leading to restrictions on water use, engine emissions, and pesticide applications. Although the weekend ritual of firing up the lawn mower to once again do battle with the forces of nature is likely to continue for some generations to come, the pressure is on to reduce, if not altogether abandon, the crusade for the perfect lawn.

DEFORESTATION PROBLEMS

Forests cover large portions of the world. They produce oxygen and provide homes for people and wildlife, including some of the most threatened and endangered animals. Over 1.6 billion people rely on forest benefits, including food, water, clothing, traditional medicine, and shelter. Yet, many of these forests worldwide are under threat from deforestation.

Deforestation, or the removal of forests, can be caused by a variety of factors. Fires, clear-cutting for agriculture, ranching and development, unsustainable logging, and degradation



due to climate change can all lead to the loss of forests. It has been estimated that we are losing about 18.7 million acres of forests annually.

Deforestation can have substantial impacts on the environment. It can lead to the increased greenhouse gas emissions. Forests can act as carbon sinks when left intact, but they act as carbon sources when they are cut, burned, or otherwise removed. In Sumatra, for example, deep peatlands are being cleared, drained, and converted into pulp plantations. This action is contributing to global greenhouse gas emissions. Additionally, Water cycles can be disrupted by the removal of forests. Trees play a key role in the water cycle, helping balance water between land and the atmosphere. Deforestation can also lead to increased soil erosion as tree roots play an important role anchoring fertile soil. Finally, deforestation disrupts the lives of both humans and other animals. Millions of people depend on the forests, using it for hunting, gathering, and medicine. But deforestation impacts all of these peoples. Further, animals count on these areas as key habitat, providing them with space, food, and water.

LOSS OF BIODIVERSITY

Plant communities provide food, help purify water, generate oxygen, and supply raw materials (building, clothing, paper, etc.), yet these communities are under threat from development, invasive species, climate change, and other factors.

Endangered Species

Various factors, including human activities and climatic changes, have led to the reduction and alteration in plant populations. In response, governments and public groups have formed groups which encourage and commission studies on rare and endangered plants or plants of unknown status. The International Union for Conservation of Nature (IUCN) tries to monitor and report on both plant and animal populations worldwide. The IUCN Red List of Threatened Species is a world-renowned database of information collected over the last four decades. The IUCN Red List assesses both plants and animals and provides taxonomic, conservation status, and distribution information. The IUCN Red List sorts each species into one of the following categories:

Extinct – a species or taxon is extinct when there is no reasonable doubt that the last individual of this group has died. Exhaustive surveys of known and expected habitat during appropriate times will have failed to record the presence of this species.

Extinct in the Wild – a species is considered to be extinct in the wild when they are only known to survive in cultivation (e.g. farming), in captivity (e.g. zoo), or as a naturalized population well outside their past range. As with extinct animals, exhaustive surveys of known and expected historical habitat during appropriate times will have failed to record the presence of this species.

Critically Endangered – a species is considered to be critically endangered when all evidence indicates that its population has either: (a) been seen to be reduced by 90% or more in last 10 years or three generations, (b) its geographic range has been reduced to less than 100 km² and severely fragmented or less than 10 km², (c) population less than 250 mature individuals and continuing to decline, (d) population size of less than 50 individuals, or (e) quantitative modeling suggests the probability of extinction at least 50% in the next 10 years. It is considered to be facing an extremely high risk of extinction in the wild.

Endangered – a species is endangered when the evidence indicates that its population has either: (a) been seen to be reduced by 70% or more in last 10 years or three generations, (b) its geographic range has been reduced to less than 5000 km² and severely fragmented or less than 500 km², (c) population less than 2500 mature individuals and continuing to decline, (d) population size of less than 250 individuals, or (e) quantitative modeling suggests the probability of extinction at least 20% in the next 10 years. It is considered to be facing a very high risk of extinction in the wild.

Vulnerable – a species is considered vulnerable when its population meets any of the following criteria: (a) been seen to be reduced by 50% or more in last 10 years or three generations, (b) its geographic range has been reduced to less than 20 000 km² and severely fragmented or less than 2000 km², (c) population less than 10 000 mature individuals and continuing to decline, (d) population size of less than 1000 individuals, or (e) quantitative modeling suggests the probability of extinction at least 10% in the next 10 years. It is considered to be facing a high risk of extinction in the wild.

Near Threatened – a species that is near threatened is close to meeting the criteria for critically endangered, endangered or vulnerable in the near future.

Least Concern – a species is least concern when it does not meet any criteria to qualify for critically endangered, endangered, vulnerable, or near threatened. Species that are widespread or abundant are included in this category.



Western Prairie Fringed-Orchid

Canada has a national Species At Risk Act, the purpose of which is “to prevent wildlife species in Canada from disappearing, to provide for the recovery of wildlife species that are extirpated (no longer exist in the wild in Canada), endangered, or threatened as a result of human activity, and to manage species of special concern to prevent them from becoming endangered or threatened.” Examples of Manitoba plant species listed under this act include Rough Agalinis (Endangered) and Western Spiderwort (Threatened).

In Manitoba, we currently have eight endangered plant species. They are protected under *The Endangered Species and Ecosystems Act*. Under the act:

- It is unlawful to kill, injure, possess, disturb or interfere with the species;
- Destroy, disturb or interfere with the habitat of the species;
- Damage, destroy, obstruct or remove a natural resource on which the species depends for its life and propagation;
- Endangered or threatened ecosystems are protected.

The endangered Manitoba plant species listed under *The Endangered Species and Ecosystems Act* are: Gastony's Cliffbrake, Gattinger's Agalinis, Great Plains Ladies'-Tresses, Rough Agalinis, Smooth Goosefoot, Small White Lady's-slipper, Western Ironweed, and Western Prairie Fringed-orchid

Gastony's Cliffbrake - Gastony's Cliffbrake is listed as Endangered under Manitoba's Endangered Species and Ecosystems Act. In Manitoba, this fern species only grows on limestone rock cliffs and ledges in the Fisher Branch and Grand Rapids areas.



Gastony's Cliffbrake is an endangered fern species in Manitoba
© C. Hamel/Nature Conservancy of Canada

CLIMATE CHANGE


Climate change, or the alteration and lasting change of the distribution of weather patterns over period of time, is something that the earth is now facing. Of all the ways in which human activity affects the distribution and abundance of wildlife on our planet, none is as pervasive and powerful as climate change. All species have a capability to adapt – at least to some degree – to natural stresses. Changes to climate and habitat have been occurring for eons, and with them have come changes to the diversity of species on earth. What makes current climate change unique is that, with the exception of cataclysmic events such as

meteor strikes, the rate at which it is taking place is leaving species and ecosystems no time to adapt.

The direct impacts of human caused climate change have now been documented on every continent, in every ocean, and in most major taxonomic groups. The increase in storms and unpredictable weather patterns is also expected with climate change. These extreme weather events can devastate plant populations as well as their habitat. This puts already vulnerable species further at risk of extinction.

There have been a variety of predictions made about the impacts of climate change on plant communities:

- If species are unable to change distribution fast enough, they may be 'left behind' and die off
- Species with long life cycles or slow dispersal rates are especially vulnerable
- Species that are isolated or found in specific separated areas are vulnerable as they may have nowhere to go. Communities include:
 - Arctic and alpine species
 - Island species
 - Costal species found between human development and rising sea levels
- Some plant communities may be lost as species move and adapt at different rates
- Increase invasive species invasions may occur, as the conditions of an area may become more suitable for the invasive species compared the native species.
- Many plant communities currently act as 'sinks' and store carbon. This helps offset carbon emissions. However, the effects of climate change on these plants may mean that they may become more of a source of carbon instead of a sink in years to come.



**Estimate shifts in plant communities in a mountain region
due to temperature increases expected with climate change**

© 2014 R.G. Bailey/Springer Science

It has been estimated that plants with thicker leaves, efficient water-use strategies, deeper roots and higher wood density are better adapted to cope with future climate change.

SKILLS AND METHODS IN PLANT SCIENCE

VEGETATION CHARACTERISTICS

Vegetation structure and flower composition are often measured as a plant community. A few specific measures are taken:

Stratification – the arrangement of plants in layers

Cover – as a percentage, the surface area of the sample plant covered in plants

Phytomass – the mass of plants, expressed as a dry mass (g/m^2 or kg/m^2) or productivity (g/m^2 or kg/m^2 per year)

Leaf area index (LAI) - the projected area of leaves over a unit of land ($\text{m}^2 \text{ m}^{-2}$), so one unit of LAI is equivalent to 10 000 m^2 of leaf area per hectare

Species composition – a list of the species that can be found within a defined area (e.g., a meter squared, an acre, etc.)

Species abundance – the amount or quantity of specific species that can be found within a defined area (e.g., a meter squared, an acre, etc.)

PATHOGENS AND DISEASE MANAGEMENT

Pathogens are a natural part of any ecosystem. Species within the environment have evolved alongside parasites and pathogens. Parasites induce specific reactions and changes in the immunity, physiology, and host behaviour. The hosts, or plants, use these changes to counteract the parasite attack. The actions of parasites on populations help select for individuals with increased resistance to this parasite's invasion. Parasites also keep evolving

alongside their hosts. However, the movement of plant material around the world has increased the movement of invasive or non-native pathogens into naive forest ecosystems.

Plant pathogens can affect the whole plant, causing defoliation, root decay and stem cankers that reduce the distribution of nutrients. Pathogens can include fungi, bacteria, viruses, as well as animals such as invertebrates. Over 80-85% of the diseases of plants are caused by fungi.

Some species of fungi can attack living rather than dead organisms and as such are the most important cause of plant diseases. More than 5000 species of fungi attack agricultural crops, garden plants, trees, and many wild plants.

The same qualities that make fungi pests on plants can also make them valuable. For example, certain yeasts can produce ethanol and carbon dioxide and play an important role in baking, brewing, and winemaking. Further, certain antibiotics, including penicillin, is produced by fungi.



Late blight on potato leaf
© Howard F. Schwartz/Colorado State

Some species of fungi can attack living rather than dead organisms and as such are the most important cause of plant diseases. More than 5000 species of fungi attack agricultural crops, garden plants, trees, and many wild plants.

The same qualities that make fungi pests on plants can also make them valuable. For example, certain yeasts can produce ethanol and carbon dioxide and play an important role in baking, brewing, and winemaking. Further, certain antibiotics, including penicillin, is produced by fungi.



Late blight on tomato stem
© Scot Nelson

Additionally, many species of invertebrates can directly impact the health of a tree. Some invertebrates may defoliate trees or bore into them while going through one of the stages of their life cycle. Wood boring insects are those that bore into the wood during the larval stage. Invasive wood boring insects are often moved through the transportation of wood products into new areas, in addition to the natural movement of the species.

The spread of invasive tree pests or pathogens is of great concern to the forestry industry, as well as citizens, first nations, and anyone who is concerned about the health and well-being of our forests. Canadian and worldwide communities depend on healthy forests. The rise in invasive pathogens in our forested areas can have impacts on biodiversity, ecosystem health, human health, natural resource industries, and international trade.

The spread of Emerald Ash Borer (EAB) occurred at a much quicker rate than expected in the eastern part of North America because of the movement of firewood. Since its accidental introduction from Asia, this invasive pest has killed untold millions of ash trees in forest, riparian, and urban settings. The larval stage (grub) of EAB can stay in the wood and exits upon arrival to a new area. The challenge with wood boring insects is that they grow within the tree and can be difficult to detect until they have already created damage and potential tree mortality. Unchecked, EAB could functionally extirpate ash with devastating economic and ecological impacts.

Foliage feeding insects eat the leaves of trees, and multiple years of defoliation can cause tree mortality. These insects have a native range in which they can move, however humans speed up the movement of these species by transporting a life stage of that insect. For example, the Gypsy Moth can be introduced to new forests through the movement of the larval stage of the insect (the caterpillar) by a car, boat or other means to a new ecosystem. The larva or caterpillar of the Gypsy Moth is the damaging stage as it consumes the leaves of trees. They can consume tremendous amounts of leaf material in a short period of time (as much as one square foot of leaves per day). When populations reach outbreak proportions, the caterpillars can completely defoliate host trees over a wide geographic area. Consistent or repeated defoliation over several years can have devastating effects, often leading to tree stress and death.

Galls – Galls are a swelling growth that can be found on the external tissues of plants (see below). Plant galls are a response by plants to an alien substance in their tissues, such as fungi, bacteria, insects, or mites. For example, a parasitic egg of an insect can cause the plants cells to transform into a gall.



FORESTRY

Forestry Management

Manitoba's vision of environmentally sound and sustainable economic growth and forestry is governed by the following principles and guidelines (these guidelines have equal status to the principles).

Conservation – Maintain essential ecological processes, biological diversity and life-support systems of our environment; harvest renewable resources on a sustained-yield basis; and make wise and efficient use of our renewable and non-renewable resources.

Enhancement – Enhance the long-term productive capability, quality and capacity of our natural ecosystems.

Global Responsibility – This principle requires that we think globally when we act locally. There is a need to work cooperatively within Canada, and internationally, to accelerate the merger of environment and economics in decision making and to develop comprehensive and equitable solutions to problems.

Integration of Environmental and Economic Decisions – Ensure that economic decisions adequately reflect environmental impacts including that on human health. Environmental initiatives shall adequately take into account economic consequences.

Prevention – Anticipate, prevent or mitigate significant adverse environmental (including human health) and economic impacts of policy, programs and decisions.

Recycling – Reduce, reuse and recover the products of our society.

Rehabilitation and Reclamation – Rehabilitation and reclamation require repairing damage caused in the past. Future policies, programs and developments should take into consideration the need for rehabilitation and reclamation.

Scientific and Technological Innovation - To research, develop, test and implement technologies essential to further environmental quality including human health and economic growth.

Shared Responsibility – Acknowledge responsibility for sustaining the environment and the economy, with each being accountable for decisions and actions, in a spirit of partnership and open cooperation.

Stewardship – Stewardship requires the recognition that we are caretakers of the environment and the economy for the benefit of present and future generations of Manitobans. A balance must be struck between today's decisions and tomorrow's impacts.

Forest Ecosystem Based Management

Forest Ecosystem Based Management is the process of developing management principles and implementing actions (measures for that forest ecosystem that will preserve and ensure its stability and sustainability). The greater the biodiversity of a forest ecosystem the more stable it is and vice versa. (An ecosystem is a self-sustaining and independent interaction of abiotic and biotic factors in a community where these interactions occur.)

We are all part of natural ecosystems, and any study or management of those ecosystems must take us into account. In spite of our culturally constructed economic and social systems, they still depend on and are very much involved in the natural cycles of ecosystems.

In order to develop an ecosystem approach, they must take an ecosystem point of view, which means a total systems approach and includes many things omitted in a less comprehensive view, i.e. no longer a single-purpose approach to the environment (harvesting trees without regard for wildlife, water, etc.). An ecosystem approach requires we take into account the relationship between artificial and natural environments and people.

In any given forest area, the forest manager has available not one but many different harvesting practices that can be adapted to the site, i.e. the kind of forest, its location, the site conditions, the species composition, and management objectives. Today, timber harvests have to be planned around the timber, the structure and function of the forest ecosystem, the wildlife it holds, and esthetics. Based on social demands, the forest manager can adapt or modify the harvest, as the situation requires.

Forestry Measuring Techniques


Forests are measured for a variety of reasons:

- Forest growth, stand development and health monitoring
- Site productivity measurements


- Permanent sample plots (PSP) using repeated measurements on same trees over time allow growth calculations,
- Calculated volume for harvest (TSP, PHS)
- Calculation of forest regeneration status and survival (Regen and FTG)
- Dendrochronology – Tree-Ring databases, long-term climate studies, biotic and abiotic factors
- Standardized measurements using easy to use and carry equipment often with mathematical basis incorporated – ie Pythagoras, or using easy to calculate constants, ie milhectares

Diameter Measurements (Tapes, Calipers, Rulers):

Top view of tree trunk cross-section



Tape wrapped around tree



Technique for Diameter Tape:

- Use side of tape which reads: CIRC. TO DIA. π mm
- Wrap tape around tree, at standard height above ground (1.3 meters = Breast Height)
- Ensure it is horizontal and no obstructions on stem like branches, etc are affecting it
- Pull tight, read number below 0, this is the diameter, from above example = 4.3 cm (Note: Tape is graduated into tenths of centimeters)

AGRICULTURE

Agriculture is the cultivation of land and breeding of animals and plants to provide food, and other products, like fibre and medicinal plants, to sustain and enhance human life.

Sustainable agriculture

Many benefits have come with modern farming practices, including power, water and communication utilities in rural households, the ability of most citizens to pursue leisure activities and general prosperity in both rural and urban segments of society. Advances in agriculture have also reduced many health risks to farm workers as well as production and economic risks by lowering the frequency of crop failures. Current levels of wealth generation, economic productivity and overall societal progress would not be possible without the use of inputs such as fertilizers and pesticides. However, there have also been significant costs. Prominent among these are topsoil depletion, groundwater contamination, the decline of family farms, continued neglect of the living and working conditions for farm laborers, increasing costs of production, and the disintegration of economic and social conditions in rural communities.

The consequences of food and fiber production affect the viability of future generations. There is evidence that ancient civilizations declined in significant part because farming practices degraded their natural resource base. Without a sufficient supply of water and healthy soil, agricultural productivity declines. Without adequate water quality, health of crops, livestock, humans and wildlife is jeopardized. Without native plant and animal diversity, ecosystems are less able to support human activity. Ongoing dependence of agriculture on non-renewable energy sources is probably not sustainable for either agriculture or society as a whole. It is critical to increase the sustainability of agricultural systems while the natural resource base is still viable.

The three pillars of sustainable agriculture

The core principle of sustainable development in general is to meet the needs of the present without compromising the needs of future generations. Sustainable agriculture, as a part of sustainable development, rests on three pillars: the economy, the environment and society. Sustainable agriculture is *economically* sustainable – it earns a profit for those whose livelihood depends on it. It is *environmentally* sustainable – it ensures that the natural environment will continue to support life and human activity. And it is *socially* sustainable – it provides a good quality of life for farmers, ranchers, farm workers and their communities.

HORTICULTURE

Horticulture is the science of the production, marketing, and use of high-value, intensively cultivated plants. These crops include annual and perennial plants, edible, and ornamental plants. Horticulturalists play a part in the development and production of a safe and secure food supply.

Horticulture combines modern and traditional technologies for the sustainable and profitable production and marketing of specialty crops. Horticulturalists are as diverse as the crops that make up the industry and the discipline. They include a wide array of individuals and groups who farm, landscape, garden, research, advise and enjoy the bounty of horticultural plants for their nourishment, health benefits and aesthetics.

REFERENCES

The document was compiled using the following references (please note – you are NOT responsible for anything in the following documents)

- "Snow and Plants." National Snow and Ice Data Center. <https://nsidc.org/cryosphere/snow/plants.html>
- Atlas of Plant Anatomy.* <<http://atlasveg.ib.usp.br/English/>>
- Bailey, R. G. (2014) Ecoregions: The Ecosystem Geography of the Ocean and Continents. Springer Science + Media, USA
- Brunner I., Herzog C., Dawes M.A., Arend M. & Sperisen C. (2015) How tree roots respond to drought. *Frontiers in Plant Science* **6**, 1–16.
- Cesareo, K. and Walker, L.K. (2018) Deforestation. *World Wildlife Fund* <<https://www.worldwildlife.org/threats/deforestation>>
- Costa A.W., Michalski G., Schauer A.J., Alexander B., Steig E.J. & Shepson P.B. (2011) Analysis of atmospheric inputs of nitrate to a temperate forest ecosystem from $\delta^{17}\text{O}$ isotope ratio measurements. *Geophysical Research Letters* **38**.
- Dabrowski S, Van Zeumeren R, McFarlaneq J, & Shaddock J. (2015) Study Guide 2016 Invasive Species: A challenge to the environment, economy, and society. North American Envirothon (2016)
- Díaz, S., Kattge, J., Cornelissen, J. H. C., et al. (2016). The global spectrum of plant form and function. *Nature*, 529(7585), 167–171.
- Edible Wild Food (2012) Eating Garlic Mustard is a win-win. Available from <<http://www.ediblewildfood.com/blog/2012/04/eating-garlic-mustard-is-a-win-win/>>
- Encarta Encyclopaedia (2000) Photosynthesis
- Harris J.G. & Harris M.W. (2001) Plant Identification Terminology: An illustrated glossary. Spring Lake Publishing, Utah, USA.
- Hogan, M.C. (Lead Author), Taub, D. R. (Topic Editor) (2010) Plant In: *Encyclopedia of Earth*. Eds. Cutler J. Cleveland.
- Keddy, P.A. (2015) Competition in plant communities. Oxford Bibliographies in Ecology.
- Körner C. (2016) Plant adaptation to cold climates. *F1000Research* **5**, 2769.
- Lukac M., Calfapietra C., Lagomarsino A. & Loreto F. (2010) Global climate change and tree nutrition: effects of elevated CO₂ and temperature. *Tree Physiology* **30**, 1209–1220.
- Mack RN, Simberloff D, Lonsdale WM, et al (2000) Biotic Invasions: Causes, epidemiology, global consequences, and control. *Ecological Applications* 10:689–710.
- Maron J.L. & Crone E. (2006) Herbivory: Effects on plant abundance, distribution and population growth. *Proceedings of the Royal Society B: Biological Sciences* **273**, 2575–2584.
- Middleton, L. (2001). Shade-tolerant flowering plants: Adaptations and horticulture implications. *Acta horticulturae* **552**: 95-102
- Moore, Randy, W. Dennis Clark, and Kingley R. Stern. (1995) *Botany*. Boston: William C. Brown Publishers
- Natural Resources Canada. (2005) Alien Forrest Pests: Context for the Canadian Forest Service's Science Program. *Government of Canada*.
- Navarro L.M., Proença V., Kaplan J.O. & Pereira H.M. (2015) Maintaining Disturbance-Dependent Habitats. In: *Rewilding European Landscapes*. (Eds H.M. Pereira & L.M. Navarro), pp. 143–167. Springer International Publishing, Cham.
- Onoda, Y., & Anten, N. P. (2011). Challenges to understand plant responses to wind. *Plant signaling & behavior*, 6(7), 1057-9.
- Pflugfelder, B. The Chemistry of Fall Colors. *Science Bob* <<https://sciencebob.com/why-do-leaves-change-color-in-the-fall/>>
- Rai P.K. (2016) Impacts of particulate matter pollution on plants: Implications for environmental biomonitoring. *Ecotoxicology and Environmental Safety* **129**, 120–136.
- Raven, P.H., Evert, R.F., and Eichhorn, S.E. (2005) Biology of Plants. *Eds. Ahr, K., Anderson, S. Weiss, V. Moscatelli, B.* W.H. Freeman and Company, New York, USA.
- Smith, R.L. and Smith, T.M. (2001) Ecology and Field Biology. *Eds. Fogarty, E., Dutton, H.. Merquillo, C., Earl, W., Burch, B. and Hitchcock, S.* Benjamin Cummings, USA

- Sytsma, K. (2018) Vascular Flora of Wisconsin – Botany 401. University of Wisconsin Madison.
- The Biology Place*. <http://www.phschool.com/science/biology_place>, Pearson Education Inc.
- Turner, M. G. (1998). Landscape ecology, living in a mosaic. In S. I. Dodson et al., (Eds.), *Ecology* (pp. 78–122). New York: Oxford University Press.
- van der Maarel E. & Franklin J. eds (2013) *Vegetation Ecology*, 2nd edn. Wiley-Blackwell, Oxford, UK.
- Virginia Institute of Marine Science. "Loss of plant diversity threatens Earth's life-support systems, experts say." *ScienceDaily* <www.sciencedaily.com/releases/2011/03/110303153116.htm>
- Willis, K.J. (2017) State of the World Plants 2017. *Eds. Willis, K.J.*, Royal Botanic Gardens <<https://stateoftheworldplants.org/2017>>
- Winner W.E. & Atkinson C.J. (1986) Absorption of air pollution by plants, and consequences for growth. *Trends in Ecology & Evolution* 1, 15–18.